



(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
**07.07.2004 Bulletin 2004/28**

(21) Application number: **03290025.0**

(22) Date of filing: **06.01.2003**

(51) Int Cl.7: **C07D 409/06, C07D 215/44,  
C07D 401/06, C07D 413/06,  
C07D 471/04, C07D 417/06,  
C07D 401/10, C07D 405/06,  
C07D 401/14, A61K 31/4704,  
A61K 31/4709, A61P 9/10,  
A61P 17/06, A61P 29/00,  
A61P 37/00**

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR  
HU IE IT LI LU MC NL PT SE SI SK TR**  
Designated Extension States:  
**AL LT LV MK RO**

(71) Applicant: **Warner-Lambert Company LLC**  
**Morris Plains, New Jersey 07950 (US)**


(72) Inventors:  
• **Kuhn, Cyrille**  
**75015 Paris (FR)**  
• **Feru, Frédéric**  
**75011 Paris (FR)**

• **Bazin, Marc**  
**94110 Arcueil (FR)**  
• **Awad, Mohamed**  
**Westerly, Rhode Island 02891 (US)**  
• **Goldstein, Steven Wayne**  
**Noank, Connecticut 06340 (US)**

(74) Representative: **Ruddock, Keith Stephen et al**  
**Pfizer Limited,**  
**European Patent Department,**  
**Ramsgate Road**  
**Sandwich, Kent CT13 9NJ (GB)**

(54) **Quinoline derivatives as CRTH2 antagonists**

(57) The invention relates to compounds of formula (I)



wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in the description, their use as medicament, pharmaceutical compositions containing them and processes for their preparation.

**Description****Field of the invention**

[0001] The invention relates to quinoline derivatives, pharmaceutical compositions containing them, processes for their preparation and their use as medicament.

**Background of the invention.**

[0002] In 1999, Nagata et al identified CRTH2 (chemoattractant receptor-homologous molecule expressed on Th2 cells), a novel G protein-coupled receptor (GPCR) belonging to the leucocyte chemoattractant receptor family.


[0003] The CRTH2 receptor is selectively expressed from a wide variety of tissues including the brain, lung and lymphoid organs in mouse (Abe *et al.*, Gene (1999) 227 (1):71-7). With respect to expression from immune system cells, it is reported that CRTH2 receptor is selectively expressed on Th2 cells, eosinophils and basophils, but not Th1 cells, B cells and NK cells in human (Nagata *et al.*, FEBS Letters (1999) 459 (2):195-9).

[0004] Bauer *et al.* (see EP1170594A2) identified Prostaglandine D<sub>2</sub> (PGD<sub>2</sub>) as the endogenous ligand of CRTH2. PGD<sub>2</sub> is released from immunologically stimulated mast cells and Th2 cells.

[0005] Interaction of CRTH2 with PGD<sub>2</sub> plays a critical role in the allergen-induced recruitment of Th2 cells in the target tissues of allergic inflammation. In addition, CRTH2 mediates PGD<sub>2</sub> dependent cell migration of blood eosinophils and basophils.


[0006] Allergic asthma and allergic rhinitis are diseases that currently affect about 10% of the population, and that number appears to be increasing (Bush, R.K.a.G., John W., Handbook of asthma and rhinitis. 1st ed. (1997), Abingdon: Blackwell Science. 270). Currently numerous classes of pharmaceutical agents are widely used to treat these diseases, for example, antihistamines, decongestants,  $\beta$ 2 agonists, anticholinergics, methylxanthines, cromolyns, corticosteroids, and leukotriene modulators. Generally however, the usefulness of these agents is limited by side effects and low efficacy. Accordingly, there is a critical medical need to identify pharmaceutically active compounds that interfere with key steps of the inflammatory and immunological processes that contribute to these disease states, and other inflammatory conditions.

[0007] EP 987251 discloses cholesteryl ester transfer protein of formula




for use in the treatment of diseases affected by low levels of HDL cholesterol and/or high levels of LDL-cholesterol and triglycerides such as atherosclerosis and cardiovascular diseases.

[0008] WO91/05549 discloses compounds of formula




for use as vasodilator, hypotensive agent, water diuretics and platelet agglutination inhibitor.

[0009] US4521607 discloses compounds of formula



for use as antihypertensives.


[0010] WO 02/079165 discloses compounds of formula



as STAT6 signaling pathway modulators.

### Summary of the invention.

[0011] The invention relates to quinoline derivatives of general formula (I)



wherein,

- R<sup>1</sup> represents H, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>2</sub>-C<sub>4</sub>)alkenyl, (C<sub>2</sub>-C<sub>4</sub>)alkynyl or (CH<sub>2</sub>)<sub>m</sub>-R<sup>1</sup>, in which

R<sup>1</sup> is selected from aromatic heterocycle, phenyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl wherein the phenyl, the heterocycle and the cycloalkyl groups are unsubstituted or substituted by one or several groups, preferably 1 to 3, selected from

- Q<sup>1</sup>, and,
- (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several, preferably 1 to 3, groups which are the same or different and which are selected from Q<sup>1</sup>, wherein Q<sup>1</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;

m is an integer selected from 0, 1 and 2;

- R<sup>2</sup> represents (C<sub>1</sub>-C<sub>4</sub>)alkyl, wherein the alkyl group may be substituted with one or several, substituents, preferably 1 to 3, selected from halogen, OR<sup>9</sup>, NR<sup>9</sup>R<sup>10</sup>, COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NHSO<sub>2</sub>R<sup>9</sup> and C(=O)(C<sub>1</sub>-C<sub>4</sub>)alkyl wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;

- R<sup>3</sup> represents (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl or -A-R<sup>3</sup>, wherein

- A represents a bond, straight or branched (C<sub>1</sub>-C<sub>3</sub>)alkylene, or (C<sub>2</sub>-C<sub>3</sub>)alkenylene;
- R<sup>3</sup> represents (C<sub>6</sub>-C<sub>12</sub>)aryle or an heterocycle, optionally aromatic, having from 5 to 10 atoms in the cycle, wherein the aryle and the heterocycle groups are unsubstituted or substituted by one or several substituents, preferably 1 to 3, selected from,
  - an aryle, an aromatic heterocycle,
  - Q<sup>2</sup>, and,
  - (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several, preferably 1 to 3, groups which are the same or different and which are selected from Q<sup>2</sup>, wherein Q<sup>2</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, SR<sup>9</sup>, COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;
- R<sup>4</sup> represents H or (C<sub>1</sub>-C<sub>4</sub>)alkyl;
- R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are the same or different and are selected from

- H, Q<sup>3</sup>, and,
- (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several groups, preferably 1 to 3, which are the same or different and which are selected from Q<sup>3</sup>, wherein Q<sup>3</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, SR<sup>9</sup>, COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl; their optical isomers, as well as their N-oxides and pharmaceutically acceptable salts and their use as medicament.

[0012] The invention further concerns the use of a compound of formula (I) for the preparation of a medicament for the prevention or the treatment of disorders for which therapy by a CRTH2 antagonist is relevant.


[0013] The invention also provides a method for the treatment of a disorder for which therapy by a CRTH2 antagonist is relevant, comprising administering to a mammal in need thereof an effective amount of a compound of formula (I).

[0014] The invention also concerns a pharmaceutical composition comprising a compound of formula (I) together with a pharmaceutically acceptable carrier, excipient, diluent or delivery system.

[0015] The invention further relates to processes for the preparation of compounds of formula (I).

#### Detailed description of the invention.

[0016] The invention relates to quinoline derivatives of formula (I)



wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in the summary of the invention.

their optical isomers, as well as their N-oxides and pharmaceutically acceptable salts and their use as CRTH2 antagonist and as medicament.

[0017] These compounds are selective CRTH2 antagonists. In man, CRTH2 is selectively expressed on immune system cells, and in particular on Th2 cells, eosinophils and basophils. CRTH2 plays a critical role in the recruitment of these cells. The compounds of the invention are useful for the prevention or treatment of disease states involving Th2 cells, eosinophils and/or basophils. In particular, the compounds of the invention are useful in the prevention and treatment of disease states involving inflammatory components, including, without limitation, inflammatory disorders such as rheumatoid arthritis, osteoarthritis, atherosclerosis, Crohn's disease, colitis ulcerosa, inflammatory bowel disease; disorders of the skin including psoriasis, eczema, erythema, pruritis, and acne; systemic lupus erythematosus, chronic obstructive pulmonary disease, angioedema, stroke, any disease marked by reperfusion injury, graft rejection, and autoimmune diseases, allergic diseases, such as allergic asthma, atopic dermatitis, and allergic rhinitis.



[0018] The compounds of the invention are also useful as research tools that can be used to antagonize the CRTH2 receptor. For example, to determine the signaling pathway of a PGD<sub>2</sub> mediated effect, it is essential to find out which PGD<sub>2</sub> receptor is involved in said pathway. This can be achieved only if selective antagonists are available for each of the PGD<sub>2</sub> receptors. The compounds of the invention, which are selective CRTH2 antagonists can be used for that purpose.

[0019] Preferred compounds of the invention are compounds of formula (I) wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in the summary of the invention with the exclusion of:

N-(1-benzoyl-6-bromo-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-acetanilide,  
 N-(1-benzoyl-6-chloro-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-acetanilide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-(4-methoxyphenyl)-2-methyl-propanamide,  
 N-[1-(4-fluorobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-butanamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-pentanamide,  
 N-[1-[(4-fluorophenyl)acetyl]-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-propanamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-2,2-dimethyl-N-phenyl-propanamide,  
 N-(1-benzoyl-6-bromo-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-phenyl-pentanamide,  
 N-[1-(2-furanylcarbonyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 2-methyl-N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-propanamide,  
 2,2,2-trifluoro-N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-acetamide,

N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinoliny)-N-(3-methoxyphenyl)-acetamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinoliny)-N-(4-methylphenyl)-acetamide,  
 N-[1-(4-chloro-3-nitrobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinoliny]-N-phenyl-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(3-nitrobenzoyl)-4-quinoliny]-acetamide,  
 5 N-phenyl-N-[1,2,3,4-tetrahydro-1-[3-(4-methoxyphenyl)-1-oxo-2-propenyl]-2-methyl-4-quinoliny]-acetamide,  
 N-[1-(3-chlorobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinoliny]-N-phenyl-acetamide,  
 N-[1-(3-fluorobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinoliny]-N-phenyl-acetamide,  
 N-[1-[4-(1,1-dimethylethyl)benzoyl]-1,2,3,4-tetrahydro-2-methyl-4-quinoliny]-N-phenyl-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(1-oxo-3-phenyl-2-propenyl)-4-quinoliny]-acetamide,  
 10 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(2-thienylcarbonyl)-4-quinoliny]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-(4-methoxybenzoyl)-2-methyl-4-quinoliny]-acetamide,  
 N-[1-(3,5-dinitrobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinoliny]-N-phenyl-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(4-nitrobenzoyl)-4-quinoliny]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-(2-iodobenzoyl)-2-methyl-4-quinoliny]-acetamide,  
 15 N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinoliny]-acetamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinoliny)-N-phenyl-pentanamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinoliny)-N-phenyl-butanamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinoliny)-N-phenyl-propanamide,  
 1-benzoyl-1,2,3,4-tetrahydro-4-(N-phenylacetamido)-quinaldine,  
 20 N-[(1-benzoyl-1, 2, 3, 4-tetrahydro-2-methyl-4-quinoliny)-2-methyl-N-phenyl propanamide;  
 N-[1-(4-bromobenzoyl)-1,2,3,4-tetrahydro-2,6-dimethyl-4-quinoliny]-acetamide;  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2,6-dimethyl-4-quinoliny)-acetamide; and,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinoliny)-acetamide.

[0020] A further preferred group of compounds are cis-isomers of formula (Ia) and (Ib),





wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in compound of formula (I) the summary of the invention and R<sup>4</sup>  
 is (C<sub>1</sub>-C<sub>4</sub>)alkyl,  
 with the exclusion of the following compounds,

N-[(2R, 4S)-1-benzoyl-1, 2, 3, 4-tetrahydro-2-methyl-4-quinoliny]-2-methyl-N-phenyl propanamide;  
 N-[(2R, 4S)-1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinoliny]-2,2-dimethyl-N-phenyl-propanamide;  
 50 N-[(2R, 4S)-1-benzoyl-1, 2, 3, 4-tetrahydro-2-methyl-4-quinoliny]-N-phenyl butanamide;  
 N-[(2R, 4S)-1-benzoyl-1, 2, 3, 4-tetrahydro-2-methyl-4-quinoliny]-N-phenyl acetamide,  
 N-[(2R, 4S)-1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinoliny]-N-phenyl-pentanamide and,  
 N-[(2R, 4S)-1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinoliny]-acetamide.

[0021] Another preferred group of compounds are cis-isomers of formula (Ia) and (Ib),  
 wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in the summary of the invention and R<sup>4</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl with the  
 exclusion of the cis-isomers of the following compounds,

N-(1-benzoyl-6-bromo-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-acetanilide,  
 N-(1-benzoyl-6-chloro-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-acetanilide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-(4-methoxyphenyl)-2-methyl-propanamide,  
 N-[1-(4-fluorobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-butanamide,  
 5 N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-pentanamide,  
 N-[1-(4-fluorophenyl)acetyl]-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-propanamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-2,2-dimethyl-N-phenyl-propanamide,  
 N-(1-benzoyl-6-bromo-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-phenyl-pentanamide,  
 N-[1-(2-furanylcarbonyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 10 2-methyl-N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-propanamide,  
 2,2,2-trifluoro-N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-acetamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-(3-methoxyphenyl)-acetamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-(4-methylphenyl)-acetamide,  
 N-[1-(4-chloro-3-nitrobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 15 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(3-nitrobenzoyl)-4-quinolyl]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-[3-(4-methoxyphenyl)-1-oxo-2-propenyl]-2-methyl-4-quinolyl]-acetamide,  
 N-[1-(3-chlorobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 N-[1-(3-fluorobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 N-[1-(4-(1,1-dimethylethyl)benzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 20 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(1-oxo-3-phenyl-2-propenyl)-4-quinolyl]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(2-thienylcarbonyl)-4-quinolyl]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-(4-methoxybenzoyl)-2-methyl-4-quinolyl]-acetamide,  
 N-[1-(3,5-dinitrobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(4-nitrobenzoyl)-4-quinolyl]-acetamide,  
 25 N-phenyl-N-[1,2,3,4-tetrahydro-1-(2-iodobenzoyl)-2-methyl-4-quinolyl]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-acetamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-phenyl-pentanamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-phenyl-butanamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-phenyl-propanamide, 1-benzoyl-1,2,3,4-tetrahydro-  
 30 4-(N-phenylacetamido)-quinaldine,  
 N-[1-benzoyl-1, 2, 3, 4-tetrahydro-2-methyl-4-quinolyl]-2-methyl-N-phenyl propanamide;  
 N-[1-(4-bromobenzoyl)-1,2,3,4-tetrahydro-2,6-dimethyl-4-quinolyl]-acetamide;  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2,6-dimethyl-4-quinolyl)-acetamide; and,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-acetamide.

[0022] Another preferred group of compound according to the invention are trans isomers of formula (Ic) or (Id):




wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined in compound of formula (I) in the summary of the invention and  $R^4$  is  $(C_1-C_4)$ alkyl.

[0023] A preferred group of trans-isomers are those of formula (Ic) or (Id) wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined in compound of formula (I) in the summary of the invention and  $R^4$  is  $(C_1-C_4)$ alkyl with the exclusion of the

trans-isomers of the following compounds,

N-(1-benzoyl-6-bromo-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-acetanilide,  
 N-(1-benzoyl-6-chloro-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-acetanilide,  
 5 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-(4-methoxyphenyl)-2-methyl-propanamide,  
 N-[1-(4-fluorobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-butanamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-pentanamide,  
 N-[1-[(4-fluorophenyl)acetyl]-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-propanamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-2,2-dimethyl-N-phenyl-propanamide,  
 10 N-(1-benzoyl-6-bromo-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-phenyl-pentanamide,  
 N-[1-(2-furanylcabonyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 2-methyl-N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-propanamide,  
 2,2,2-trifluoro-N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-acetamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-(3-methoxyphenyl)-acetamide,  
 15 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-(4-methylphenyl)-acetamide,  
 N-[1-(4-chloro-3-nitrobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(3-nitrobenzoyl)-4-quinolyl]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-[3-(4-methoxyphenyl)-1-oxo-2-propenyl]-2-methyl-4-quinolyl]-acetamide,  
 N-[1-(3-chlorobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 20 N-[1-(3-fluorobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 N-[1-[4-(1,1-dimethylethyl)benzoyl]-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(1-oxo-3-phenyl-2-propenyl)-4-quinolyl]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(2-thienylcarbonyl)-4-quinolyl]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-(4-methoxybenzoyl)-2-methyl-4-quinolyl]-acetamide,  
 25 N-[1-(3,5-dinitrobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(4-nitrobenzoyl)-4-quinolyl]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-(2-iodobenzoyl)-2-methyl-4-quinolyl]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-acetamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-phenyl-pentanamide,  
 30 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-phenyl-butanamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-phenyl-propanamide,  
 1-benzoyl-1,2,3,4-tetrahydro-4-(N-phenylacetamido)-quinaldine,  
 N-[1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-2-methyl-N-phenyl propanamide;  
 N-[1-(4-bromobenzoyl)-1,2,3,4-tetrahydro-2,6-dimethyl-4-quinolyl]-acetamide;  
 35 N-(1-benzoyl-1,2,3,4-tetrahydro-2,6-dimethyl-4-quinolyl)-acetamide; and,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-acetamide.

[0024] Compounds of formula (I) wherein R<sup>4</sup> represents (C<sub>1</sub>-C<sub>4</sub>)alkyl, have two chiral centers, numbered C\*<sup>1</sup> and C\*<sup>2</sup> on the following formula




As each of C\*<sup>1</sup> and C\*<sup>2</sup> can have a Cahn-Ingold-prelog configuration R or S, compounds of formula (I) exist in four different forms:




C\*<sup>1</sup>(R)-C\*<sup>2</sup>(S) - (CIS),  
 C\*<sup>1</sup>(S)-C\*<sup>2</sup>(R) - (CIS),  
 C\*<sup>1</sup>(R)-C\*<sup>2</sup>(R) (TRANS), and,  
 C\*<sup>1</sup>(S)-C\*<sup>2</sup>(S) (TRANS).


[0025] As mentioned above, a compound of formula (I), wherein R<sup>4</sup> represents (C<sub>1</sub>-C<sub>4</sub>)alkyl, includes four diastereoisomers that can be represented by the following formula (Ia), (Ib), (Ic) and (Id):




(Ia)



(Ib)



(Ic)



(Id)

wherein (Ia) and (Ib) are the cis isomers and (Ic) and (Id) are the trans isomers.

[0026] In the present application, the terms "optical isomer" include the diastereoisomers of formula (Ia), (Ib), (Ic) and (Id) and mixture thereof.

[0027] Preferred compounds of formula (I) are compounds of formula (Ia), (Ib) and mixture thereof.

[0028] In the above groups of compounds, the following substitutions are further preferred:

[0029] Preferably, R<sup>1</sup> represents H, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>2</sub>-C<sub>4</sub>)alkenyl, (C<sub>2</sub>-C<sub>4</sub>)alkynyl or (CH<sub>2</sub>)<sub>m</sub>-R<sup>1</sup>, wherein

R<sup>1</sup> is selected from phenyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl wherein the phenyl and the cycloalkyl groups are unsubstituted or substituted by one or several groups, preferably 1 to 3, selected from

- Q<sup>1</sup>, and,
- (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several, preferably 1 to 3, groups which are the same or different and which are selected from Q<sup>1</sup>,

wherein Q<sup>1</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H

and (C<sub>1</sub>-C<sub>4</sub>)alkyl;

m is an integer selected from 0, 1 and 2;

and R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in the summary of the invention.

**[0030]** Preferably, R<sup>1</sup> represents (CH<sub>2</sub>)<sub>m</sub>—R<sup>1</sup>, wherein

R<sup>1</sup> is selected from phenyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl wherein the phenyl and the cycloalkyl groups are unsubstituted or substituted by 1 to 3 groups selected from OR<sup>9</sup>, COOR<sup>9</sup> and (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with a group COOR<sup>9</sup> wherein R<sup>9</sup> is selected from H or (C<sub>1</sub>-C<sub>4</sub>)alkyl,

m is an integer selected from 0 and 1;

and R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in the summary of the invention.

**[0031]** Another preferred group of compounds include compounds of formula (I) wherein R<sup>1</sup> represents (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>2</sub>-C<sub>4</sub>)alkenyl, (C<sub>2</sub>-C<sub>4</sub>)alkynyl or (CH<sub>2</sub>)<sub>m</sub>—R<sup>1</sup>, in which

R<sup>1</sup> is selected from aromatic heterocycle and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl wherein the heterocycle and the cycloalkyl groups are unsubstituted or substituted by one or several groups, preferably 1 to 3, selected from

- Q<sup>1</sup>, and,
- (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several, preferably 1 to 3, groups which are the same or different and which are selected from Q<sup>1</sup>,

wherein Q<sup>1</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;

m is an integer selected from 0, 1 and 2;

and R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in the summary of the invention.

**[0032]** Preferably R<sup>1</sup> represents a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl wherein the cycloalkyl group is unsubstituted or substituted by one or several groups, preferably 1 to 3, selected from

- Q<sup>1</sup>, and,
- (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several, preferably 1 to 3, groups which are the same or different and which are selected from Q<sup>1</sup>, wherein Q<sup>1</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;

m is an integer selected from 0, 1 and 2.

**[0033]** Preferably, R<sup>1</sup> represents a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, more preferably, a cyclopropyl group.

**[0034]** Preferably, R<sup>1</sup> is phenyl unsubstituted or substituted in the para position by a substituent selected from halogen, OR<sup>9</sup>, CH<sub>2</sub>COOR<sup>9</sup> and CH<sub>2</sub>COOR<sup>9</sup> wherein R<sup>9</sup> is selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl.

**[0035]** Preferably, R<sup>2</sup> represents (C<sub>1</sub>-C<sub>4</sub>)alkyl, optionally substituted with COOR<sup>9</sup> wherein R<sup>9</sup> is selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl.

**[0036]** Preferably, R<sup>2</sup> represents unsubstituted (C<sub>1</sub>-C<sub>4</sub>)alkyl, more preferably a methyl group.

**[0037]** R<sup>3</sup> is preferably selected from (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl and -A-R<sup>3</sup>, wherein

- A represents a bond, (C<sub>1</sub>-C<sub>3</sub>)alkylene, straight or branched, or (C<sub>2</sub>-C<sub>3</sub>)alkenylene;
- R<sup>3</sup> represents an heterocycle, optionally aromatic, having from 6 to 10 atoms in the cycle, unsubstituted or substituted by one or several, preferably 1 to 3, substituents selected from,
- an aryle, an heterocycle,
- Q<sup>2</sup>, and,
- (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several, preferably 1 to 3, groups which are the same or different and which are selected from Q<sup>2</sup>,

wherein Q<sup>2</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, SR<sup>9</sup>, COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl, with the proviso that R<sup>3</sup> is not selected from unsubstituted thienyl or unsubstituted furanyl.

[0038] Preferably, R<sup>3</sup> is selected from (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl and -A-R<sup>3</sup>, wherein

- A represents a bond, (C<sub>1</sub>-C<sub>3</sub>)alkylene, straight or branched, or (C<sub>2</sub>-C<sub>3</sub>)alkenylene;
- R<sup>3</sup> represents isoxazolyl, pyrazolyl, pyrrolyl, thiazolyl, thiadiazolyl, isothiazolyl, thiazolyl, tetrazolyl or an hetero-  
cycle, optionally aromatic, having from 6 to 10 atoms in the cycle, unsubstituted or substituted by one or several,  
preferably 1 to 3, substituents selected from,
- an aryle, an heterocycle,
- Q<sup>2</sup>, and,
- (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several, preferably 1 to 3, groups which are the same or different  
and which are selected from Q<sup>2</sup>,

wherein Q<sup>2</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, SR<sup>9</sup> COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;

[0039] Preferably, R<sup>3</sup> is selected from -A-R<sup>3</sup>, wherein

- A represents a bond;
- R<sup>3</sup> represents isoxazolyl, pyrazolyl, pyrrolyl, thiazolyl, thiadiazolyl, isothiazolyl, thiazolyl, tetrazolyl or an hetero-  
cycle, optionally aromatic, having from 6 to 10 atoms in the cycle, unsubstituted or substituted by one or several,  
preferably 1 to 3, substituents selected from,
- an aryle, an heterocycle,
- Q<sup>2</sup>, and,
- (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several, preferably 1 to 3, groups which are the same or different  
and which are selected from Q<sup>2</sup>,

wherein Q<sup>2</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, SR<sup>9</sup> COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl.

[0040] Preferably, R<sup>3</sup> is selected from -A-R<sup>3</sup>, wherein

- A represents a bond, straight or branched (C<sub>1</sub>-C<sub>3</sub>)alkylene, or (C<sub>2</sub>-C<sub>3</sub>)alkenylene;
- R<sup>3</sup> represents an aryle, preferably phenyl, unsubstituted or substituted by one or several, preferably 1 to 3, sub-  
stituents selected from,
- an aryle, an heterocycle,
- Q<sup>2</sup>, and,
- (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several, preferably 1 to 3, groups which are the same or different  
and which are selected from Q<sup>2</sup>,

wherein Q<sup>2</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, SR<sup>9</sup> COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl.

[0041] Preferably, R<sup>3</sup> is selected from -A-R<sup>3</sup>, wherein

- A represents a bond, straight or branched (C<sub>1</sub>-C<sub>3</sub>)alkylene, or (C<sub>2</sub>-C<sub>3</sub>)alkenylene;
- R<sup>3</sup> represents a phenyl, unsubstituted or substituted by 1 to 3, substituents selected from aryle, heterocycle,  
halogen, CN, CF<sub>3</sub>, OR<sup>9</sup> and COOR<sup>9</sup>. wherein R<sup>9</sup> is selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl.

[0042] Preferably, R<sup>4</sup> represents (C<sub>1</sub>-C<sub>4</sub>)alkyl, more preferably a methyl group.

[0043] Preferably R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are the same or different and are selected from H, halogen and OR<sup>9</sup> wherein R<sup>9</sup> is selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl.

[0044] In the following and in the foregoing text:

Halogen includes fluoro, chloro, bromo, and iodo.

Alkyl groups include straight or branched carbon chains. Examples of (C<sub>1</sub>-C<sub>4</sub>)alkyl groups are methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl and tert-butyl.

The term alkenyl groups include straight and branched hydrocarbon radicals having at least one double bond. Examples of (C<sub>2</sub>-C<sub>4</sub>)alkenyl groups are ethenyl, 3-buten-1-yl and the like.

The term alkynyl groups includes straight and branched hydrocarbon radicals having at least one triple bond.

Examples of (C<sub>2</sub>-C<sub>4</sub>) alkynyl are ethynyl, 3-buten-1-yl, propynyl, and the like.

The term (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl includes cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

The term heterocycle includes carbocycles, optionally aromatic, having from 5 to 10 atoms in the cycle, containing from 1 to 4 heteroatoms selected from O, S and N in the cycle. A preferred group of heterocycles includes 5 or 6-membered heterocycles containing from 1 to 3 heteroatoms selected from O, S and N.

Particularly preferred heterocycles are isoxazolyl, oxazolyl thienyl, pyrazolyl, pyrrolyl, triazolyl, tetrazolyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridinyl, pyrazinyl, benzo-oxadiazolyl or pyrazolo-pyridinyl, quinolinyl and quinoxalinyl. Aryl is understood to refer to an aromatic carbocycle containing between 6 and 12 carbon atoms. A preferred aryl group is phenyl.

[0045] Preferred compounds are:

Cis-N-[2-Methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(1-oxy-pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(4-Methoxy-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(4-Hydroxy-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(4-trifluoromethyl-benzoyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(3-phenyl-acryloyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(4-Cyano-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(4-Chloro-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 4-[Cis-4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-benzoic acid methyl ester;  
 4-[Cis-4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-benzoic acid;  
 Cis-N-[2-Methyl-1-(3-phenyl-propionyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(5-methyl-thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(Benzofurazan-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-phenylacetyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(pyrazine-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(6-Chloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(6-trifluoromethyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2,6-Dimethoxy-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2-Methoxy-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(2-methylsulfanyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2-Chloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(5-methyl-pyrazine-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2-Chloro-6-methyl-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(4-Chloro-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(6-(2,2,2-trifluoro-ethoxy)-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(2-propylsulfanyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(5,6-Dichloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2,6-Dichloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(4-methyl-[1,2,3]thiadiazole-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(5-methyl-isoxazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2,5-Dimethyl-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(1-methyl-1H-pyrrole-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(Isoxazole-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(5-methyl-isoxazole-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2,4-Dimethyl-thiazole-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(5-Chloro-thiophene-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(1,5-Dimethyl-1H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(4-methyl-isothiazole-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-5-[4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-thiophene-2-carboxylic acid dimethylamide  
 Cis-N-[1-(4-Hydroxy-quinoline-6-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(4-tert-Butyl-thiazole-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(2-Ethyl-pyridine-4-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(3,6-Dichloro-pyridine-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(4-Chloro-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-2-[4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-isonicotinic acid methyl ester  
 5 Cis-N-[2-Methyl-1-(4-[1,2,4]triazol-4-yl-benzoyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(2,6-Dimethoxy-pyridine-4-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-Ethyl-isoxazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(2-tetrazol-1-yl-pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(5-propyl-isoxazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 10 Cis-N-[1-(5-Isobutyl-2-methyl-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-Bromo-furan-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(6-phenyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(2-phenyl-pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 15 Cis-N-[2-Methyl-1-(quinoline-6-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(3,4-Dimethoxy-furan-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(3-methyl-furan-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(2,5-Dimethyl-furan-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(2,4-Dimethyl-oxazole-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 20 Cis-N-[1-(5-Methoxymethyl-furan-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-Fluoro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(quinoline-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(6-methyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(quinoline-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 25 Cis-N-[2-Methyl-1-(1H-pyrazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(2H-pyrazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-Isobutyl-isoxazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(quinoline-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(6-methyl-pyridine-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 30 Cis-N-[2-Methyl-1-(quinoxaline-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(3-Methoxy-thiophene-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-tert-Butyl-2-methyl-furan-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-Ethyl-2-methyl-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 35 Cis-N-[2-Methyl-1-([1,2,5]thiadiazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(2-methyl-5-propyl-2H-pyrazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-Benzyl-acetamide;  
 Cis-N-Benzyl-N-[2-methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 40 Trans-N-Benzyl-N-[2-methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-Cyclohexyl-N-[2-methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-(1-Benzoyl-6-methoxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 Cis-N-(1-Benzoyl-6-hydroxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 Cis-N-(1-Benzoyl-6-chloro-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 45 Cis-N-(1-Benzoyl-7-chloro-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-prop-2-ynyl-acetamide;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-(4-methoxy-phenyl)-acetamide;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-(4-hydroxy-phenyl)-acetamide;  
 Cis-[4-[Acetyl-(1-benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-amino]-phenyl]-acetic acid ethyl ester;  
 50 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-malonamic acid methyl ester;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-malonamic acid;  
 Cis-N-[2-Methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-Cyclopropyl-acetamide;  
 Cis-N-Cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide  
 55 (+)-Cis-N-cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide  
 (-)-Cis-N-cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-Cyclopropyl-N-[2-methyl-1-(3-methyl-isoxazole-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-Phenyl-N-[1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;

Cis-N-(1-Benzoyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide  
 Cis-N-[2-Ethyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-Ethyl-N-[2-Methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide.

5 [0046] The following compounds are particularly preferred:

Cis-N-[2-Methyl-1-(pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(1-oxy-pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(4-Hydroxy-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 10 Cis-N-[2-Methyl-1-(4-trifluoromethyl-benzoyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(4-Cyano-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(4-Chloro-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 4-[Cis-4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-benzoic acid methyl ester;  
 4-[Cis-4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-benzoic acid;  
 15 Cis-N-[2-Methyl-1-(3-phenyl-propionyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(5-methyl-thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(Benzofurazan-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-(2-Methyl-1-phenylacetyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(pyrazine-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 20 Cis-N-[1-(6-Chloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(6-trifluoromethyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2,6-Dimethoxy-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2-Methoxy-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(2-methylsulfanyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 25 Cis-N-[1-(2-Chloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(5-methyl-pyrazine-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2-Chloro-6-methyl-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(4-Chloro-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 30 Cis-N-[2-Methyl-1-(6-(2,2,2-trifluoro-ethoxy)-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(2-propylsulfanyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(5,6-Dichloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2,6-Dichloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 35 Cis-N-[2-Methyl-1-(4-methyl-[1,2,3]thiadiazole-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(5-methyl-isoxazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2,5-Dimethyl-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(1-methyl-1H-pyrrole-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(isoxazole-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 40 Cis-N-[2-Methyl-1-(5-methyl-isoxazole-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2,4-Dimethyl-thiazole-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(5-Chloro-thiophene-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(1,5-Dimethyl-1H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(4-methyl-isothiazole-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 45 Cis-5-[4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-thiophene-2-carboxylic acid dimethylamide  
 Cis-N-[1-(4-Hydroxy-quinoline-6-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(4-tert-Butyl-thiazole-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(2-Ethyl-pyridine-4-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 50 Cis-N-[1-(3,6-Dichloro-pyridine-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(4-Chloro-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-2-[4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-isonicotinic acid methyl ester  
 Cis-2-[4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-isonicotinic acid methyl ester  
 Cis-N-[2-Methyl-1-(4-[1,2,4]triazol-4-yl-benzoyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 55 Cis-N-[1-(2,6-Dimethoxy-pyridine-4-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-Ethyl-isoxazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(2-tetrazol-1-yl-pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(5-propyl-isoxazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(5-Isobutyl-2-methyl-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-Bromo-furan-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(6-phenyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 5 Cis-N-[2-Methyl-1-(2-phenyl-pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(quinoline-6-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(3,4-Dimethoxy-furan-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(3-methyl-furan-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(2,5-Dimethyl-furan-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 10 Cis-N-[1-(2,4-Dimethyl-oxazole-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-Methoxymethyl-furan-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-Fluoro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(quinoline-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(6-methyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 15 Cis-N-[2-Methyl-1-(quinoline-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(1H-pyrazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(2H-pyrazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-Isobutyl-isoxazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(quinoline-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 20 Cis-N-[2-Methyl-1-(6-methyl-pyridine-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(quinoxaline-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(3-Methoxy-thiophene-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-tert-Butyl-2-methyl-furan-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-Ethyl-2-methyl-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 25 Cis-N-[2-Methyl-1-([1,2,5]thiadiazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(2-methyl-5-propyl-2H-pyrazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-Benzyl-acetamide;  
 30 Cis-N-Benzyl-N-[2-methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Trans-N-Benzyl-N-[2-methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-Cyclohexyl-N-[2-methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-(1-Benzoyl-6-methoxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 Cis-N-(1-Benzoyl-6-hydroxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 35 Cis-N-(1-Benzoyl-6-chloro-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-prop-2-ynyl-acetamide;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-(4-methoxy-phenyl)-acetamide;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-(4-hydroxy-phenyl)-acetamide;  
 Cis-[4-[Acetyl-(1-benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-amino]-phenyl]-acetic acid ethyl ester;  
 40 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-malonamic acid methyl ester;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-malonamic acid;  
 Cis-N-[2-Methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-Cyclopropyl-acetamide;  
 Cis-N-Cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide  
 45 (+)-Cis-N-cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide  
 (-)-Cis-N-cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-Cyclopropyl-N-[2-methyl-1-(3-methyl-isoxazole-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-Phenyl-N-[1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-(1-Benzoyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide  
 50 Cis-N-[2-Ethyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-Ethyl-N-[2-Methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-yl]-acetamide;

**[0047]** The above compounds of configuration C\*<sup>1</sup>(R)-C\*<sup>2</sup>(S), C\*<sup>1</sup>(S)-C\*<sup>2</sup>(R) or mixture thereof are particularly preferred:

**[0048]** Most preferred compounds are:


Cis-N-[2-Methyl-1-(pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(4-Hydroxy-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[2-Methyl-1-(4-trifluoromethyl-benzoyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(4-Chloro-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(5-methyl-thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(Benzofurazan-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 5 Cis-N-[1-(6-Chloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(6-trifluoromethyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2,6-Dimethoxy-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(6-(2,2,2-trifluoro-ethoxy)-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-  
 acetamide;  
 10 Cis-N-[2-Methyl-1-(5-methyl-isoxazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(Isioxazole-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(5-Chloro-thiophene-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(5-propyl-isoxazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(quinoline-6-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 15 Cis-N-[1-(5-Methoxymethyl-furan-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-Isobutyl-isoxazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(3-Methoxy-thiophene-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-([1,2,5]thiadiazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-Benzyl-N-[2-methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 20 Cis-N-(1-Benzoyl-6-hydroxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 Cis-N-(1-Benzoyl-6-chloro-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-prop-2-ynyl-acetamide;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-(4-methoxy-phenyl)-acetamide;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-(4-hydroxy-phenyl)-acetamide;  
 25 Cis-[4-[Acetyl-(1-benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-amino]-phenyl]-acetic acid ethyl ester;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-malonamic acid methyl ester;  
 Cis-N-[2-Methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-Cyclopropyl-acetamide;  
 Cis-N-Cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide  
 30 (+)-Cis-N-cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide  
 (-)-Cis-N-cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-Cyclopropyl-N-[2-methyl-1-(3-methyl-isoxazole-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-Phenyl-N-[1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide, and  
 Cis-N-[2-Ethyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide.

[0049] The invention further relates to the above compounds or groups of compounds as medicaments. The invention further relates to the above compounds or groups of compounds as CRTH2 antagonists.

#### General process for the preparation of compounds of the invention


[0050] The invention also relates to a process for manufacturing a compound of formula (I),






wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in the summary of the invention and R<sup>4</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl, said process comprising the following steps:

(1) reacting a compound of formula (1a),



**(1a)**


wherein R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in the summary of the invention, with sodium methoxide, in the presence of a solvent such as methanol, to give the compound of formula (1b),



**(1b)**

in which R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above;


(2) reacting said compound of formula (1b) with a compound of formula R<sup>4</sup>-MgCl, wherein R<sup>4</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl in an anhydrous ether such as tetrahydrofuran and then with benzylchloroformate, to give the compound of formula (1c),



**(1c)**


in which R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above;

(3) reacting said compound of formula (1c) with a compound of formula R<sup>1</sup>-NH<sub>2</sub>, wherein R<sup>1</sup> is as defined in the summary of the invention, in the presence of a base and of TiCl<sub>4</sub> to give the compound of formula (1d),




in which R<sup>1</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above;

20 (4) reducing said compound of formula (1d), using for example NaBH<sub>4</sub> and MeOH in the presence of acetic acid, to give the compound of formula (1e),




40 in which R<sup>1</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above;

45 (5) reacting said compound of formula (1e) with a compound of formula R<sup>2</sup>COCl, wherein R<sup>2</sup> is as defined in the summary of the invention, in the presence of a base, to give the compound of formula (1f),



20 in which R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above;

(6) deprotecting said compound of formula (1f) by hydrogenolysis, using for example ammonium formate catalyzed by palladium on charcoal, to give the compound of formula (1g)



40 in which R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above;



(7) reacting said compound of formula (1g) with R<sup>3</sup>COCl, wherein R<sup>3</sup> is as defined in the summary of the invention, in the presence of a base, to give a compound of formula (I) wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above R<sup>4</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl,

45 (8) isolating said compound of formula (I), or,



(9) separating compounds of formula (Ia) and (Ib)

50

55




from compounds of formula (Ic) and (Id)




and,

- isolating said compounds of formula (Ia) and (Ib) and compounds of formula (Ic) and (Id), or
- further separating compounds of formula (Ia) and (Ib) and compounds of formula (Ic) and (Id) to obtain four separated diastereoisomers and isolating them.

**[0051]** The invention also relates to a process for manufacturing a compound of formula (Ia) or (Ib) or a mixture thereof,




(Ia)




(Ib)

wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> and R<sup>4</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl, said process being characterized in that a compound of formula (1d)



(1d)

wherein R<sup>1</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in the summary of the invention and R<sup>4</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl, is reacted with a mineral borohydride such as sodium triacetoxy borohydride in acetic acid, to give a compound cis-(1e) of formula





Cis-(1e)

wherein  $R^1$ ,  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined above.


Steps 5, 6 and 7 are then performed as disclosed in scheme 1 to give a mixture of compound of formula (Ia) and (Ib) wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined above. Optionally, compounds of formula (Ia) and (Ib) may be separated and isolated.

[0052] The invention also relates to a process for manufacturing a compound of formula (Ia) or (Ib) or a mixture thereof,



said process being comprising the following steps,


(1) reacting a compound of formula (3a)



wherein  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined in the summary of the invention, with


- a compound of formula  $R^4-MgCl$ , wherein  $R^4$  is  $(C_1-C_4)$ alkyl in an anhydrous ether such as tetrahydrofuran, and then with,
- a compound of formula  $R^3-COCl$ ,

to give a compound of formula (3b)




wherein R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above;

(2) reacting said compound of formula (3b) with a compound of formula R<sup>1</sup>-NH<sub>2</sub>, wherein R<sup>1</sup> is as defined in the summary of the invention, in the presence of a base to give a compound of formula (3c),



wherein R<sup>1</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above;

(3) reducing said compound of formula (3c) using for example a mineral borohydride such as sodium triacetoxy borohydride in acetic acid to give a compound of formula (3d),





wherein R<sup>1</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above;

(4) reacting said compound of formula (3d) with a compound of formula R<sup>2</sup>COCl, wherein R<sup>2</sup> is as defined in the summary of the invention, in the presence of a base, to give a mixture of compounds of formula (1a) and (1b) wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above;

(5) isolating said mixture, or


(6) separating and isolating said compounds of formula (1a) and (1b).

[0053] The invention also relates to a process for manufacturing a compound of formula (1a) or (1b) or mixture thereof,




said process being comprising the following steps,

(1) reacting a compound of formula (4a)




with a compound of formula  $R^1-NH_2$ , wherein  $R^1$  is as defined in the summary of the invention, to give the compound of formula (4b),



in which  $R^1$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined above;


(2) reacting said compound of formula (4b), with a compound of formula  $R^2COCl$ , wherein  $R^2$  is as defined in the summary of the invention, in the presence of a base, to give the compound of formula (4c),






in which R<sup>1</sup>, R<sup>2</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above;

(3) reacting said compound of formula (4c) with benzyl-halogen in a solvent such as acetone to give the compound of formula (4d),




in which R<sup>1</sup>, R<sup>2</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above;

(4) optionally, reacting said compound of formula (4d) with a compound of formula R<sup>4</sup>-MgCl, wherein R<sup>4</sup> is (C<sub>1</sub>-C<sub>4</sub>) alkyl in an anhydrous ether such as tetrahydrofuran to give the compound of formula (4e),




in which R<sup>1</sup>, R<sup>2</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above and R<sup>4</sup> is (C<sub>1</sub>-C<sub>4</sub>) alkyl;

(5) reducing said compound of formula (4d) or (4e), for example by hydrogenation in MeOH in presence (R,R)-(-)-1,2-bis[(o-methoxyphenyl)(phenyl)phosphino]ethane (1,5-cyclooctadiene)rhodium (I) tetrafluoroborate or NiCl<sub>2</sub>/NaBH<sub>4</sub>, to give a compound of formula (4f)



in which R<sup>1</sup>, R<sup>2</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above and R<sup>4</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>) alkyl;

(6) deprotecting said compound of formula (4f) by hydrogenolysis using for example ammonium formate catalysed by palladium on charcoal, to give a compound of formula (4g)



in which  $R^1$ ,  $R^2$ ,  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined above;

(7) reacting said compound of formula (4g) with  $R^3\text{COCl}$ , wherein  $R^3$  is as defined in the summary of the invention, in the presence of a base, to give a mixture compounds of formula (la) and (lb) wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined above;

(8) isolating said compounds of formula (la) and (lb), or,

(9) separating compounds of formula (la) and (lb) and isolating them.

[0054] Depending of their substitution, compounds of formula (I) may have further chiral centers. Optical isomers due to said further chiral centers are included within the scope of the invention.

[0055] The compounds utilized in the invention include pharmaceutically acceptable derivatives of compounds of formula (I) such as solvates, hydrates, pharmaceutically acceptable salts and polymorphs (different crystalline lattice descriptors).

[0056] Pharmaceutically acceptable salts of a compound of formula (I) include salts having a basic part and salts having an acidic part.

[0057] The expression pharmaceutically acceptable salt of a compound of formula (I) having a basic part should be understood to refer to the addition salts of the compounds of formula (I) which may be formed from non-toxic inorganic or organic acids such as, for example, hydrobromic, hydrochloric, sulfuric, phosphoric, nitric, acetic, succinic, tartaric, citric, maleic, hydroxymaleic, benzoic, fumaric and toluenesulfonic acid salts, and the like. The various quaternary ammonium salts of the derivatives (I) are also included in this category of compounds of the invention. In addition, the expression pharmaceutically acceptable salt of a compound of formula (I) having an acidic part is understood to refer to the usual salts of the compounds of formula (I) which may be formed from non-toxic inorganic or organic bases such as, for example, the hydroxides of alkali metals and alkaline-earth metals (sodium, potassium, magnesium and calcium), amines (dibenzylethylenediamine, trimethylamine, piperidine, pyrrolidine, benzylamine and the like) or alternatively quaternary ammonium hydroxides such as tetramethylammonium hydroxide. (See also "Pharmaceutical salts" by Berge S.M. *et al.* (1997) *J. Pharm. Sci.* **66**: 1-19, which is incorporated herein by reference.).

[0058] Use of a prodrug of a compound of the invention such as it would occur to one skilled in the art (see Bundgaard, *et al.*, *Acta Pharm. Suec.*, 1987; **24**: 233-246), is also contemplated.

#### Pharmaceutical compositions.

[0059] The products of the invention are administered in the form of compositions, which are appropriate for the nature, and severity of the complaint to be treated. The daily dose in humans is usually between 1 mg and 1 g of product, which may be taken in one or more individual doses. The compositions are prepared in forms which are compatible with the intended route of administration, such as, for example, tablets, coated tablets, capsules, mouthwashes, aerosols, powders for inhalation, suppositories, enemas, foams (such as rectal foams), gels or suspensions. These compositions are prepared by methods which are familiar to those skilled in the art and comprise from 0.5 to 60% by weight of active principle (compound of the invention) and 40 to 99.5% by weight of a pharmaceutical vehicle or carrier which is appropriate and compatible with the active principle and the physical form of the intended composition.

[0060] Solid form preparations include powders, tablets, dispersible granules, capsules, cachets, and suppositories.

A solid carrier can be one or more substances which may also act as diluents, flavouring agents, solubilizers, lubricants, suspending agents, binders, or tablet disintegrating agents; it can also be an encapsulating material. In powders, the carrier is a finely divided solid, which is in a mixture with the finely divided active component. In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired. The powders, tablets, cachets or encapsulated forms for capsules preferably contain 5% to about 70% of the active component. Suitable carriers are magnesium carbonate, magnesium stearate, talc, lactose, sugar, pectin, dextrin, starch, tragacanth, methyl cellulose, sodium carboxymethyl cellulose, a low-melting wax, cocoa butter, and the like.

[0061] Tablets, powders, cachets, and capsules can be used as solid dosage forms suitable for oral administration. The drug may be delivered as a spray (either in a pressurized container fitted with an appropriate valve or in a non-pressurized container fitted with a metering valve).

[0062] Liquid form preparations include solutions, suspensions, and emulsions.

[0063] Sterile water or water-propylene glycol solutions of the active compounds may be mentioned as an example of liquid preparations suitable for parenteral administration. Liquid preparations can also be formulated in solution in aqueous polyethylene glycol solution.

[0064] Aqueous solutions for oral administration can be prepared by dissolving the active component in water and adding suitable colorants, flavouring agents, stabilizers, and thickening agents as desired. Aqueous suspensions for oral use can be made by dispersing the finely divided active component in water together with a viscous material such as natural synthetic gums, resins, methyl cellulose, sodium carboxymethyl cellulose, and other suspending agents known to the pharmaceutical formulation art.

[0065] For preparing suppository preparations, a low-melting wax such as a mixture of fatty acid glycerides and cocoa butter is first melted and the active ingredient is dispersed therein by, for example, stirring. The molten homogeneous mixture is then poured into convenient sized molds and allowed to cool and solidify. Enemas are obtained according to known procedures to prepare solutions adapted for rectal administration. Foams are prepared according to known methods (these foams can notably be similar to those used to administer a drug such as 5-ASA for treating rectocolite).

[0066] Preferably the pharmaceutical preparation is in unit dosage form. In such form, the preparation is divided into unit doses containing appropriate quantities of drug. The unit dosage form can be a packaged preparation, the package containing discrete quantities of the preparation, for example, packaged tablets, capsules, and powders in vials or ampoules. The unit dosage form can also be a capsule, cachet, or tablet itself, or it can be the appropriate number of any of these packaged forms.

### Use

[0067] The compounds of the invention are CRTH2 antagonists, preferably selective CRTH2 antagonists. These compounds have low  $IC_{50}$  values, typically at most  $10\mu M$ , preferably below  $1\mu M$ , more preferably below  $500nM$ .

[0068] Compounds of the invention can be used in the prevention and the treatment of disease that can be treated by a CRTH2 antagonist such as Th2 cells-related diseases, eosinophils-related diseases and basophils-related diseases. Preferably, the compounds of the invention can be used for the prevention or the treatment of diseases involving inflammatory components including, without limitation, inflammatory disorders such as rheumatoid arthritis, osteoarthritis, atherosclerosis, Crohn's disease, colitis ulcerosa, inflammatory bowel disease; disorders of the skin including psoriasis, eczema, erythema, pruritis, and acne; systemic lupus erythematosus, chronic obstructive pulmonary disease, angioedema, stroke, and any disease marked by reperfusion injury, graft rejection, and autoimmune diseases, allergic diseases, such as allergic asthma, atopic dermatitis, and allergic rhinitis.

[0069] The invention finally relates to a method for the treatment of the above-mentioned diseases comprising administering to a mammal, particularly a human, in need thereof an effective amount of compound of the invention.


### Processes for synthesizing compounds of the invention

#### Protocol A

[0070] Protocol A is useful for the preparation of compounds of formula (I) wherein  $R^4$  is  $(C_1-C_4)$  alkyl.

[0071] Scheme 1 illustrates the preparation of cis isomers of compounds of formula (I) wherein  $R^4$  is  $(C_1-C_4)$  alkyl.

Scheme 1



[0072] The starting compounds are either commercially available or can be prepared according to routes known to the skilled person.

In step 1, compound (1a), where R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in the summary of the invention, is reacted with sodium methoxide (preferably 5-10 eq) in methanol under reflux to form the desired methoxy quinoline.

In step 2, compound (1b) is reacted with an excess of a solution of R<sup>4</sup>-magnesium chloride in anhydrous ether (for example tetrahydrofuran) at low temperature, preferably between 0 and -78°C, and then treated with an excess of benzyl chloroformate in the same condition of temperature to give compound (1c).

In step 3, compound (1c) is reacted with an amine (for example aniline), a solution of titanium chloride (preferably

1 to 1.1 eq) in toluene, a base (preferably 4 to 8 eq) such as triethyl amine, in a chlorinated solvent such as dichloromethane at a temperature below 0°C to give compound (1d).

In step 4, compound (1d) is reacted with an excess of sodium triacetoxymethyl borohydride in acetic acid at room temperature to give the desired amine (cis).

In step 5, compound cis-(1e) is acylated with an acyl chloride of formula R<sup>2</sup>-COCl (preferably 5-30 eq) in a solvent such as dioxane or DMF and a base such as diisopropylethyl amine under reflux to give compound cis-(1f).


In step 6, compound cis-(1f) is deprotected by hydrogenolysis (for example ammonium formate catalyzed by palladium on charcoal) in an alcohol, such as ethanol, under reflux, to form compound cis-(1g).

In step 7, compound cis-(1g) is reacted with an acyl chloride (commercially available or prepared from the corresponding acid) of formula R<sup>3</sup>-COCl, in a solvent such as dioxane, THF or DMF, and a base in solution or its solid supported form, such as diisopropylethyl amine, at room temperature to give a compound of formula (I) wherein R<sup>4</sup> is (C<sup>1</sup>-C<sup>4</sup> alkyl).

[0073] Optionally, cis-isomers of configuration C<sup>\*1</sup>(R) - C<sup>\*2</sup>(S) or C<sup>\*1</sup>(S) - C<sup>\*2</sup>(R) can be separated by using methods known to the skilled man, such as flash chromatography.

[0074] Scheme 2 illustrates the preparation of a mixture of diastereoisomers of compounds of formula (I) wherein R<sup>4</sup> is (C<sub>1</sub>-C<sub>4</sub>) alkyl, starting from compound (1d) of scheme 1.

Scheme 2




[0075] Compound 1h is a mixture of 4 diastereoisomers, containing two isomers with a cis configuration and two

isomers with a trans configuration. Cis isomers and trans isomers can be separated using methods known to the skilled man, such as flash chromatography.

### Protocole B

[0076] Protocol B is an alternative process for the preparation of compounds of formula (I) wherein  $R^4$  is ( $C_1$ - $C_4$ ) alkyl. Scheme 3 illustrates the preparation of cis-isomers of compounds of formula (I).

Scheme 3



[0077] Compound (3a) is obtained according to step 1 of scheme 1.

In step 1, compound (3b) is reacted with a solution of  $R^4$ -MgCl (preferably 5 to 10 eq) in anhydrous ether such as tetrahydrofuran at low temperature, preferably between  $0^\circ\text{C}$  and  $-78^\circ\text{C}$ , and then treated with  $R^3$ -COCl (preferably 5 to 10 eq) in the same condition of temperature to form compound (3b).

In step 2, compound (3b) is reacted with a compound of formula  $R^1$ -NH<sub>2</sub>, (preferably 2 to 3 eq.), titanium chloride (preferably 1 eq) in solution in toluene, TEA (preferably 4 to 8 eq.) in a chlorinated solvent, such as dichloromethane or dichloroethane at a temperature below  $0^\circ\text{C}$  and stirred over night at a temperature preferably comprised between room temperature and  $80^\circ\text{C}$ , to give the imine derivative (3c).

In step 3, compound (3c) is reduced with a mineral borohydride (preferably 5 to 6 eq) such as sodium triacetoxyborohydride, in acetic acid. The reaction is carried out at room temperature to form the amine (3d).

In step 4, compound (3d) is acylated with  $R^2$ -COCl (preferably 1 to 6 eq), in a solvent such as dioxane or DMF, and a base such diisopropylethyl amine (preferably 1 to 3 eq) at room temperature to give the desired compound of formula (I).


[0078] A mixture of diastereoisomers of compounds of formula (I) can be obtained by replacing step 3 of scheme 3 by step 4 of scheme 2. The trans isomers can then be obtained using classical methods of diastereoisomers separation.

### Protocole C

[0079] Protocol C is an alternative process for the preparation of compounds of formula (I) wherein  $R^4$  is ( $C_1$ - $C_4$ ) alkyl or hydrogen.

[0080] Scheme 4 illustrates the preparation of cis isomers of compounds of formula (I) wherein  $R^4$  is H or ( $C_1$ - $C_4$ ) alkyl.

Scheme 4



In step 1, compounds (4a) is reacted with a compound of formula  $R^1-NH_2$ , optionally in a solvent such as acetonitrile, preferably at a temperature between  $150^\circ C$  to  $200^\circ C$  for 0.5 to 1 hour, to form compound (4b).

In step 2, compound (4b) is acylated with an acid chloride of formula  $R^2-COCl$  (preferably 5 to 10 eq), in an appropriate solvent such as pyridine or dioxane with or without N-ethyl-N,N-diisopropylamine, preferably at a temperature of between  $50$  and  $130^\circ C$  to form compound (4c).

In step 3, compound (4c) is reacted with benzyle-halogen, said halogen being preferably Cl or Br, in a solvent such as acetone to obtain compound of formula (4d).

Step 4 is performed to obtain compounds of formula (I) in which  $R^4$  is  $(C_1-C_4)$  alkyl. Step 4 is as disclosed in Scheme 1. Alternatively, to obtain compounds of formula (I) where  $R^4$  is H, step 5 is performed directly after step 3. In step 5, compound (4d) (from step 3) or compound (4e) (from step 4) is reduced into compound (4f). The reduction is preferably carried out using  $NiCl_2/NaBH_4$  in MeOH/THF or by hydrogenation in MeOH in presence (R,R)-(-)-1,2-bis[(o-methoxyphenyl)(phenyl)phosphino]ethane(1,5-cyclooctadiene)rhodium (I) tetrafluoroborate.

In step 6, compound (4f) is deprotected by hydrogenolysis (for example ammonium formate catalysed by palladium



on charcoal) in an alcohol, such as ethanol, under reflux, to form compound (4g).

In step 7, compound (4g) acylated with a compound of formula  $R^3\text{-COCl}$  (preferably 5 to 10 eq), in an appropriate solvent such as pyridine or dioxane with or without N-ethyl-N,N-diisopropylamine, preferably at a temperature of between 50 and 130°C to form compound (4h).

### Synthesis Examples

[0081] The following examples illustrate, without limiting it, the synthesis of particularly active compounds of formula (I) according to the invention.

[0082]  $^1\text{H}$ -NMR spectra were recorded at 400 MHz. The residual solvent peak, usually chloroform ( $\delta_{\text{H}}$  7.27 ppm) or DMSO ( $\delta_{\text{H}}$  2.5 ppm) was used as internal shift reference. Analytical HPLC was run on a DIONEX Summit equipped with a diode array detector, using a Kromasil C-18 reversed-phase column and eluting with the following general system: acetonitrile with 0.1% v/v of acid formic in water with 0.1 % v/v of acid formic (5:95 to 95:5 in a 13 minutes gradient). LC/MS studies were run on a Hewlett-Packard HPLC 1100 coupled to a Micromass mass spectrometer.

[0083] HPLC: the purity was obtained at 214 nm, and expressed as a percentage of areas (are of the considered peak vs total of the peak areas).

### Preparation of intermediates

#### Intermediate A

##### *Cis-N-(2-Methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide*

[0084] Intermediate A corresponds to a compound of formula cis-(1g), prepared according to step 1 to 6 of scheme 1 ( $R^1$ : Phenyl,  $R^2$ :  $\text{CH}_3$ ,  $R^4$ :  $\text{CH}_3$ ,  $R^5$ : H,  $R^6$ : H,  $R^7$ : H,  $R^8$ : H).

##### 4-Methoxy Quinoline (compound of formula (1b))

[0085] To a solution of sodium methoxide (33g) in methanol (130ml), 4-chloroquinoline (10g) was added under stirring. The mixture was heated under reflux for 6 hours.

[0086] After cooling, the reaction mixture was concentrated, then dissolved in dichloromethane (250ml) and washed with water (250ml). The organic layer was dried over sodium sulfate. The solvent was evaporated to give a yellow oil (7.39g, 76% yield) which was not purified.

$^1\text{H}$  NMR [ $(\text{CD}_3)_2\text{SO}$ ]:  $\delta$  8.8 (m, 1H), 8.15 (m, 1H), 8 (m, 1H), 7.8 (m, 1H), 7.6 (m, 1H), 7 (m, 1H), 4 (s, 3H).

MS positive ESI:  $m/z$  ( $m+H$ ) $^{+}$ = 159

HPLC:  $r_T$ : 10.77, 97.16% purity

##### 2-Methyl-4-oxo-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (compound of formula (1c))

[0087] To a solution of 4-methoxy quinoline (3g) in tetrahydrofuran (80 ml), under nitrogen, and under cooling at -78°C in an acetone-dry ice bath and under stirring, a solution of methyl magnesium chloride at 3 M/l in tetrahydrofuran was added dropwise (77 ml) for 30 minutes. The mixture was stirred for 30 minutes and then benzyl chloroformate (32.15g) was added dropwise for 45 minutes at -78°C. The mixture was stirred at room temperature over night. The reaction mixture was hydrolyzed with methanol (200ml) and then with HCl 1mol/l (200 ml) and finally concentrated. Dichloromethane (200ml) and aqueous solution of sodium bicarbonate were added to the mixture under stirring. The separate organic layer was washed with an aqueous solution of sodium chloride (200ml). The solvent was removed under reduce pressure to give an oil, which was chromatographed in silica gel with an eluent of cyclohexane / ethyl acetate (95/5). The fractions containing the desired compound were combined and evaporated to give an oil (5.44g, 98% yield).

$^1\text{H}$  NMR [ $(\text{CD}_3)_2\text{SO}$ ]:  $\delta$  7.9 (m, 2H), 7.6 (t, 1H), 7.3-7.5 (m, 5H), 7.2 (m, 1H), 5.3 (d, 2H), 5.1 (t, 1H), 3.2 (m, 1H), 2.5 (m, 1H), 1.15 (d, 3H). HPLC:  $r_T$ : 7.30, 96.9% purity

##### 2-Methyl-4-phenylimino-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (compound of formula (1d))

[0088] To a solution of 2-Methyl-4-oxo-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (9g), triethylamine (24.65g), aniline (5.7g) in dichloromethane (110 ml), cooled at -5°C in an ice/sodium chloride bath, a solution of titanium chloride in toluene 1 mol/l was dropwise added for 10 minutes. The mixture was stirred over night at room temperature and then washed with dichloromethane and an aqueous solution of potassium carbonate. The organic layer was sep-

arated, dried over sodium sulfate and the solvent was removed under reduced pressure to give a brown oil (16g) which was directly used in the next step.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 8.3 (m, 1H), 7.8 (m, 1H), 7.3-7.5 (m, 8H), 7.2 (m, 1H), 7.1 (m, 1H), 6.8 (m, 2H), 5.2 (dd, 2H), 5 (m, 1H), 2.75 (m, 1H), 2.55 (m, 1H), 1.15 (m, 3H).

#### Cis-2-methyl-4-phenylamino-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (Compound of formula cis-(1e))

**[0089]** To a solution of 2-methyl-4-phenylimino-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (16g) in acetic acid (450ml), sodium triacetoxyborohydride was added in small amounts (32.32g), under stirring, at room temperature. The reaction mixture was then stirred at room temperature for one hour and poured on 1 liter of cold water under stirring. After precipitation, the mixture was filtered, dissolved in ethyl acetate and then, washed with water. The organic layer was separated and dried over sodium sulfate. The solvent was removed to give the crude compound (9g), recrystallised in hot cyclohexane (40ml), and then filtered at room temperature to give the desired compound (5.3g, 46.6% yield).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.5 (m, 1H), 7.15-5.5 (m, 9H), 7.1 (m, 1H), 6.75 (m, 1H), 6.65 (m, 2H), 5.25 (dd, 2H), 4.6 (m, 1H), 4.3 (m, 1H), 3.8 (m, 1H), 2.6 (m, 1H), 1.4 (m, 1H), 1.25 (m, 3H).

HPLC: *r*<sub>T</sub>: 12.50, 100% purity

#### Cis-4-(acetyl-phenylamino)-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (compound of formula cis-(1f))

**[0090]** To a solution of cis-2-methyl-4-phenylamino-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (5.3 g), diisopropyl ethyl amine (2.73g) in dioxane (31ml), anhydride acetic (7.26g) was added under stirring. The mixture was heated under reflux over night. After cooling, dichloromethane and an aqueous solution of potassium carbonate were added. The organic layer was separated, and then dried over sodium sulfate. Organic solvent was removed under reduce pressure to give the crude compound (18g). The desired compound (3.8g, 64% yield) was obtained after purification by flash chromatography on silica gel with cyclohexane/ethyl acetate (80/20) as eluent and then cyclohexane/ethyl acetate (50/50).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.1-7.5 (m, 14H), 5.45 (m, 1H), 5.2 (dd, 2H), 4.45 (m, 1H), 2.2 (m, 1H), 2.05 (s, 3H), 1.3 (m, 1H), 1.1 (m, 3H).

HPLC: *r*<sub>T</sub>: 11.64, 100% purity

#### Intermediate A (compound of formula cis-(1g))

**[0091]** To a solution of cis-4-(acetyl-phenylamino)-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (3.8g), ammonium formate (5.7g) in ethanol (90ml), palladium (PD/C at 10% few milligrams) was added. The reaction mixture was heated under reflux for two hours. After cooling the mixture was filtered on celite and the solvent was evaporated under reduce pressure to give intermediate A (2.3g, 90% yield).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.25 (m, 4H), 7.0 (m, 3H), 6.75 (t, 1H), 6.45 (d, 1H), 6.3 (m, 1H), 3.55 (m, 2H), 1.9 (m+s, 4H), 1.35 (m, 1H), 1.1 (d, 3H).

HPLC: *r*<sub>T</sub>: 8.14, 100% purity

#### Intermediate B

##### ***N*-Benzyl-(2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-acetamide**

**[0092]** Intermediate B corresponds to compound of formula (1g) wherein R<sup>1</sup>: benzyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H and R<sup>8</sup>: H.

Intermediate Cis-B (Cis-*N*-Benzyl-(2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-acetamide) and Trans-B (Trans-*N*-Benzyl-(2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-acetamide) were prepared as follows:

#### 2-Methyl-4-benzylimino-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (compound of formula (1d))

**[0093]** To a solution of 2-Methyl-4-oxo-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (1.00 g, 3.39 mmol), triethylamine (3.77 mL, 27.1 mmol) in dichloromethane (10 mL), cooled at 0°C in an ice/sodium chloride bath, a solution of titanium tetrachloride in toluene (3.39 mL, 1.0 M in toluene) was dropwise added for 10 minutes. Benzyl amine (0.74 mL, 3.39 mmol) was added and the mixture was stirred over night at room temperature. The reaction was quenched with saturated NaHCO<sub>3</sub> solution (100 mL) and the mixture was extracted with ethyl acetate (3 x 200 mL). The combined

organic layers were washed with brine (200 mL), dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated to give a reddish oil (1.70 g) which was used directly in the next step.

$^1\text{H}$  NMR [ $\text{CDCl}_3$ ]:  $\delta$  8.3 (d, 1H), 7.6 (d, 1H), 7.1-7.5 (m, 12H), 5.3 (d, 1H), 5.2 (d, 1H), 5.1 (m, 1H), 4.7 (d, 1H), 4.6 (d, 1H), 2.8 (m, 2H), 1.1 (d, 3H).

MS positive ESI:  $m/z$  ( $m+H$ ) $^+$  = 385

Cis and Trans-2-Methyl-4-benzylamino-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (Compound of formula (1e))

[0094] To a solution of 2-Methyl-4-benzylamino-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (1.70 g, 3.38 mmol) in methanol (50 mL), was added sodium borohydride (2.56 g, 67.7 mmol) in small portions. The reaction was stirred overnight and then concentrated to near dryness under vacuum. This was then diluted with a concentrated solution of potassium carbonate (50 mL) and then extracted with ethyl acetate (3 x 150 mL). The combined organic layers were washed with brine (200 mL), dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated to give a pale yellow oil (1.34 g).

This material was purified on silica gel with 15% ethyl acetate / hexane to provide two fractions:

Higher R<sub>f</sub> fraction - Cis-2-Methyl-4-benzylamino-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (300 mg):  $^1\text{H}$  NMR [ $\text{D}_6$  DMSO]:  $\delta$  7.5 (d, 1H), 7.4 (d, 2H), 7.1-7.3 (m, 11H), 5.1 (q, 1H), 5.2 (d, 2H), 4.3 (m, 1H), 3.9 (m, 2H), 2.8 (m, 1H), 1.1 (d, 3H), 1.0 (m, 1H).

MS positive ESI:  $m/z$  ( $m+H$ ) $^+$  = 387

[0095] A portion of this material was treated with HCl gas and recrystallized to provide a single crystal which was identified as the cis diastereomer by X-ray diffraction.

[0096] Lower R<sub>f</sub> fraction — Trans-2-Methyl-4-benzylamino-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (300 mg):  $^1\text{H}$  NMR [ $\text{D}_6$  DMSO]:  $\delta$  7.5 (d, 1H), 7.4-7.2 (m, 8H), 7.0 (t, 1H), 5.2 (q, 2H), 4.5 (m, 1H), 3.7 (m, 1H), 3.6 (m, 2H), 2.5 (m, 1H), 2.3 (m, 1H), 1.5 (m, 1H), 1.1 (d, 3H).

MS positive ESI:  $m/z$  ( $m+H$ ) $^+$  = 387

Cis-4-(acetyl-benzylamino)-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (compound of formula cis-(1f))

[0097] To a solution of Cis-2-Methyl-4-benzylamino-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (300 mg, 0.776 mmol), diisopropyl ethyl amine (0.135 mL, 0.776 mmol) in dioxane (1.5 mL), was added anhydride acetic (0.732 mL, 7.76 mmol) with stirring. The mixture was heated under reflux over night, cooled and directly chromatographed on silica gel utilizing 50% ethyl acetate / hexane to provide the title compound (300 mg) as an oil.  $^1\text{H}$  NMR [ $\text{CDCl}_3$ ]:  $\delta$  7.6-6.9 (m, 14H), 5.5 (d, 1H), 5.3 (d, 1H), 5.2 (t, 1H), 4.8 (m, 1H), 4.4 (m, 2H), 3.8 (d, 1H), 2.4 (s, 3H), 1.5 (m, 1H), 1.2 (m, 3H).

MS positive ESI:  $m/z$  ( $m+\text{NH}_4$ ) $^+$  = 446

[0098] Trans-4-(acetyl-benzylamino)-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (compound of formula trans-(1f)) When the Trans isomer (300 mg) was utilized in the above reaction, the title compound (320 mg) was isolated as an oil.  $^1\text{H}$  NMR [ $\text{CDCl}_3$ ]:  $\delta$  7.8 (d, 1H), 7.4-7.0 (m, 13H), 6.6 (m, 1H), 5.2 (s, 2H), 4.8 (m, 1H), 4.4 (d, 2H), 4.1 (d, 1H), 2.4, 2.1 (s, 3H), 2.0 (m, 1H), 1.2 (d, 3H), 1.3 (m, 2H).

MS positive ESI:  $m/z$  ( $m+\text{NH}_4$ ) $^+$  = 446

**Cis-*N*-Benzyl-(2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-acetamide (intermediate Cis-B)**

[0099] To a solution of Cis-4-(acetyl-benzylamino)-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid benzyl ester (290 mg, 0.677 mmol) and methanol (10 mL) was added 10% Pd-C (100 mg). Hydrogen gas was then added via a balloon, and the solution was stirred rapidly. After 1h, the reaction was filtered, the residual catalyst washed with toluene, and the organic layers combined and concentrated to give the title compound as an oil.  $^1\text{H}$  NMR [ $\text{CDCl}_3$ ]:  $\delta$  7.4-7.2 (m, 5H), 7.1 (m, 1H), 6.9 (m, 1H), 6.7 (m, 1H), 6.5 (m, 1H), 6.3 (br s, 1H), 5.3, 5.1 (m, 1H), 4.5 (d, 1H), 4.2 (d, 1H), 3.9-3.5 (m, 3H), 2.4, 2.1 (s, 3H), 2.1, 1.9 (m, 1H), 1.5 (m, 1H), 1.1 (m, 3H).

MS positive ESI:  $m/z$  ( $m+H$ ) $^+$  = 295

**Trans-*N*-Benzyl-(2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-acetamide (intermediate Trans-B)**

Trans-4-(acetyl-benzylamino)-2-methyl-3,4-dihydro-2H-quinoline

[0100] When the trans isomer (306 mg) was utilized in the above reaction, the title compound was isolated as an oil.  $^1\text{H}$  NMR [ $\text{CDCl}_3$ ]:  $\delta$  7.4-7.2 (m, 3H), 7.2 (d, 1H), 7.1 (t, 1H), 6.9 (d, 1H), 6.7 (m, 2H), 6.0 (t, 1H), 5.1, 4.9 (m, 1H),

4.5 (d, 1H), 4.4 (d, 1H), 3.9-3.3 (m, 3H), 2.4, 2.1 (s, 3H), 2.1, 1.9 (m, 1H), 1.5 (m, 2H), 1.1 (m, 3H).  
MS: 295, MH<sup>+</sup>

### Example 1

#### Cis-N-[2-Methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

[0101] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: thien-2-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

[0102] To a solution of intermediate A (0.1g), diisopropyl ethyl amine (0.05g) in dioxane (1ml), was added 2-thiophen-carbonyl chloride (0.057mg), under stirring, at room temperature for 3 hours. Then dichloromethane (2 ml) and water (2 ml) were added. The organic layer was separated and the solvent was removed under reduce pressure. The crude compound was crystallized in diethyl ether, filtered and dried to give a white solid (0.055g, 39.6% yield).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.4-7.2 (m, 8H), 7.05 (m, 1H), 6.85 (t, 1H), 6.75 (t, 1H), 6.7 (d, 1H), 5.5 (m, 1H), 4.7 (m, 1H), 2.3 (m, 1H), 2.0 (s, 3H), 1.6 (s, 1H), 1.1 (d, 3H).

HPLC: r<sub>T</sub>: 10.15, 100% purity

MS positive ESI: m/z (m+H)<sup>+</sup>=391

### Example 2

#### Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide

[0103] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.4-7.15 (m, 12H), 6.9 (t, 1H), 6.5 (d, 1H), 5.6 (m, 1H), 4.8 (m, 1H), 2.3 (m, 1H), 2.0 (s, 3H), 1.6 (s, 1H), 1.1 (d, 3H).

HPLC: r<sub>T</sub>: 10.15, 100% purity

MS positive ESI: m/z (m+H)<sup>+</sup>=385

### Example 3

#### Cis-N-[2-Methyl-1-(pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

[0104] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: pyridin-4-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 8.5 (d, 2H), 7.4 (m, 4H), 7.2 (m, 3H), 7.0 (d, 2H), 6.9 (t, 1H), 6.45 (d, 1H), 5.6 (m, 1H), 4.8 (m, 1H), 2.3 (m, 1H), 2.0 (s, 3H), 1.65 (s, 1H), 1.1 (d, 3H).

HPLC: r<sub>T</sub>: 7.76, 98.2% purity

MS positive ESI: m/z (m+H)<sup>+</sup>=386

### Example 4

#### Cis-N-[2-Methyl-1-(1-oxy-pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

[0105] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 1-oxy-pyridin-4-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

To a solution of Cis-N-[2-methyl-1-(pyridine-4-carbonyl)-1,2,3,4-tetrahydroquinolin-4-yl]-N-phenyl-acetamide (0.1g) in dichloromethane (2.6 ml) was added meta-chloro-para-benzoic acid (0.26mmol). After 3 hours, dichloromethane was added and the solution was washed using water and carbonate potassium. The organic layer was separated and dried over sodium sulfate. The solvent was removed to give the title compound (60 mg).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 8.0 (d, 2H), 7.4 (m, 3H), 7.2 (m, 4H), 7.0 (m, 3H), 6.5 (d, 1H), 5.55 (m, 1H), 4.7 (m, 1H), 2.3 (m, 1H), 2.0 (s, 3H), 1.15 (m+d, 4H).

HPLC: r<sub>T</sub>: 7.58, 97% purity

MS positive ESI: m/z (m+H)<sup>+</sup>=402

### Example 5

#### Cis-N-[1-(4-Methoxy-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

[0106] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 4-methoxyphenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.4 (m, 4H), 7.3 (m, 2H), 7.2 (m, 3H), 6.9 (t, 1H), 6.7 (d, 2H), 6.5 (d, 1H), 5.6 (m, 1H), 4.75 (m, 1H), 3.75 (s, 3H), 2.3 (m, 1H), 2.0 (s, 3H), 1.6 (s, 1H), 1.15 (d, 3H).

HPLC: *r*<sub>T</sub>: 10.20, 97.7% purity

MS positive ESI: *m/z* (*m*+H)<sup>+</sup>= 415

### Example 6

#### Cis-N-[1-(4-Hydroxy-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

[0107] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 4-hydroxyphenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

To a solution N-[1-(4-Methoxy-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide (0.1g) in dichloromethane (2.4 ml) at 0°C, was added boron tribromide (0.265mmol). After 3 hours, ice and ethyl acetate were added and the solution was washed using water and carbonate potassium. The organic layer was separated and dried over sodium sulfate. The solvent was removed to afford an amorphous material. The product was purified by chromatography on silica gel using a mixture of dichloromethane and ethanol (95:5) as eluent to give the title compound (50 mg).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.95 (s, 1H), 7.4-7.2 (m, 6H), 7.1 (t, 1H), 6.9 (m, 3H), 6.5 (d, 1H), 6.4 (d, 2H), 5.5 (m, 1H), 4.65 (m, 1H), 2.2 (m, 1H), 2.0 (s, 3H), 1.05 (m+d, 4H). HPLC: *r*<sub>T</sub>: 9.08, 95.6% purity

MS positive ESI: *m/z* (*m*+H)<sup>+</sup>=401

### Example 7

#### Cis-N-[2-Methyl-1-(4-trifluoromethyl-benzoyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

[0108] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 4-trifluoromethylphenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.4-7.1 (m, 11H), 6.85 (t, 1H), 6.35 (d, 1H), 5.55 (m, 1H), 4.7 (m, 1H), 2.25 (m, 1H), 2.0 (s, 3H), 1.1 (d+m, 4H).

HPLC: *r*<sub>T</sub>: 11.36, 93.6% purity

MS positive ESI: *m/z* (*m*+H)<sup>+</sup>=453

### Example 8

#### Cis-N-[2-Methyl-1-(3-phenyl-acryloyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

[0109] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: -CH=CH-phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.7 (d, 1H), 7.4-7.2 (m, 13H), 7.1 (d, 1H), 6.6 (d, 1H), 5.4 (m, 1H), 4.75 (m, 1H), 2.25 (m, 1H), 2.0 (s, 3H), 1.6 (s, 1H), 1.15 (d, 3H).

HPLC: *r*<sub>T</sub>: 10.92, 100% purity

MS positive ESI: *m/z* (*m*+H)<sup>+</sup>=411

### Example 9

#### Cis-N-[1-(4-Cyano-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

[0110] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 4-cyano-benzoyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.4-7.2 (m, 11H), 6.9 (t, 1H), 6.4 (d, 1H), 5.6 (m, 1H), 4.8 (m, 1H), 2.3 (m, 1H), 2.0 (s, 3H), 1.15 (d+m, 4H).

HPLC: *r*<sub>T</sub>: 10.05, 99.5% purity

MS positive ESI: *m/z* (*m*+H)<sup>+</sup>= 410

**Example 10****Cis-N-[1-(4-Chloro-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0111] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 4-chloro-phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.4-7.1 (m, 11H), 6.9 (t, 1H), 6.45 (d, 1H), 5.6 (m, 1H), 4.75 (m, 1H), 2.3 (m, 1H), 2.0 (s, 3H), 1.1 (d+m, 4H).

HPLC: r<sub>T</sub>: 10.98, 99.5% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 419

**Example 11****4-[cis-4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-benzoic acid methyl ester**

[0112] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 4-benzoic acid methyl ester, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.9 (d, 2H), 7.4-7.25 (m, 8H), 7.15 (t, 1H), 6.9 (m, 1H), 6.4 (d, 1H), 5.6 (m, 1H), 4.8 (m, 1H), 3.9 (s, 3H), 2.3 (m, 1H), 2.0 (s, 3H), 1.15 (d+m, 4H).

HPLC: 100% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 443

**Example 12****4-[Cis-4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-benzoic acid**

[0113] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 4-benzoic acid, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

A solution of 4-[4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-benzoic acid methyl ester (50mg) and lithium hydroxyde (0.283mmol) in methanol (2ml) and water (1ml) was stirred at room temperature overnight. After addition of water, the mixture was washed with dichloromethane. The aqueous layer was acidified and then extracted using ethyl acetate. The organic layer is separated and dried over sodium sulfate. The solvent was removed to give the title compound (50mg).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.9 (d, 2H), 7.4-7.2 (m, 9H), 6.9 (m, 1H), 6.4 (d, 1H), 5.6 (m, 1H), 4.8 (m, 1H), 2.3 (m, 1H), 2.0 (s, 3H), 1.15 (d+m, 4H).

HPLC: r<sub>T</sub>: 8.92, 100% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 429

**Example 13****Cis-N-[2-Methyl-1-(3-phenyl-propionyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0114] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: -(CH<sub>2</sub>)<sub>2</sub>-phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.4-7.2 (m, 10H), 7.1 (m, 3H), 6.95 (d, 1H), 5.2 (m, 1H), 4.7 (m, 1H), 3.0-2.6 (m, 4H), 2.1 (m, 1H), 2.0 (s, 3H), 1.0 (d+m, 4H).

HPLC: r<sub>T</sub>: 10.71, 100% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 413

**Example 14****Cis-N-[2-Methyl-1-(5-methyl-thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0115] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 5-methyl-thien-2-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

[0116] The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.4 (m, 4H), 7.25 (m, 3H), 7.1 (t, 1H), 6.9 (d, 1H), 6.5 (d, 1H), 6.4 (d, 1H), 5.5 (m, 1H), 4.6 (m, 1H), 2.3 (s, 3H), 2.2 (m, 1H), 2.0 (s, 3H), 1.1 (d+m, 4H).

HPLC: r<sub>T</sub>: 10.51, 95.3% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 405

**Example 15****Cis-N-[1-(Benzofurazan-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0117] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: benzo[1,2,5]oxadiazol-5-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.9 (s, 1H), 7.5 (m, 1H), 7.3 (m, 4H), 7.15 (m, 3H), 6.9 (m, 2H), 6.5 (m, 1H), 5.5 (m, 1H), 4.7 (m, 1H), 2.25 (m, 1H), 2.0 (s, 3H), 1.1 (d+m, 4H).

HPLC: r<sub>T</sub>: 10.92, 100% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 427

**Example 16****Cis-N-(2-Methyl-1-phenylacetyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenylacetamide**

[0118] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: benzyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.3-7.2 (m, 7H), 7.1-7.0 (m, 7H), 5.0 (m, 1H), 4.6 (m, 1H), 3.65 (dd, 2H), 2.1 (m, 1H), 1.9 (s, 3H), 1.0 (d+m, 4H).

HPLC: r<sub>T</sub>: 10.41, 100% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 399

**Example 17****Cis-N-[2-Methyl-1-(pyrazine-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0119] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: pyrazin-2-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 18****Cis-N-[1-(6-Chloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0120] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 6-chloro-pyridin-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 19****Cis-N-[2-Methyl-1-(6-trifluoromethyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0121] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 6-trifluoromethyl-pyridin-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 20****Cis-N-[1-(2,6-Dimethoxy-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0122] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 2,6-dimethoxy-pyridin-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 21****Cis-N-[1-(2-Methoxy-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0123] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 2-methoxy-pyridin-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 22**

**Cis-N-[2-Methyl-1-(2-methylsulfanyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0124] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 2-methylsulfanyl-pyridin-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 23**

**Cis-N-[1-(2-Chloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0125] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 2-chloro-pyridin-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 24**

**Cis-N-[2-Methyl-1-(5-methyl-pyrazine-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0126] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 5-methyl-pyrazin-2-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 25**

**Cis-N-[1-(2-Chloro-6-methyl-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0127] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 2-chloro-6-methyl-pyridin-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 26**

**Cis-N-[1-(4-Chloro-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0128] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 4-chloro-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridin-5-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 27**

**Cis-N-[2-Methyl-1-(6-(2,2,2-trifluoro-ethoxy)-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0129] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 6-(2,2,2-trifluoroethoxy)-pyridin-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 28**

**Cis-N-[2-Methyl-1-(2-propylsulfanyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0130] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 2-propylsulfanyl-pyridin-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.



**Example 29****Cis-N-[1-(5,6-Dichloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

- 5 [0131] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 5,6-dichloro-pyridin-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 30**

10 **Cis-N-[1-(2,6-Dichloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

- [0132] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 2,6-dichloro-pyridin-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

15 **Example 31**

**Cis-N-[2-Methyl-1-(4-methyl-[1,2,3]thiadiazole-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

- 20 [0133] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 4-methyl-[1,2,3]-thiadiazol-5-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 32**

25 **Cis-N-[2-Methyl-1-(5-methyl-isoxazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

- [0134] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 5-methyl-isoxazol-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

30 **Example 33**

**Cis-N-[1-(2,5-Dimethyl-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

- 35 [0135] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 2,5-dimethyl-2H-pyrazol-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 34**

40 **Cis-N-[2-Methyl-1-(1-methyl-1H-pyrrole-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

- [0136] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 1-methyl-1H-pyrazol-2-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

45 **Example 35**

**Cis-N-[1-(Isioxazole-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

- 50 [0137] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: isoxazol-5-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 36****Cis-N-[2-Methyl-1-(5-methyl-isoxazole-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

- 55 [0138] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 5-methyl-isoxazol-4-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 37**

**Cis-N-[1-(2,4-Dimethyl-thiazole-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

- 5    **[0139]**   R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 2,4-dimethyl-thiazol-5-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

**Example 38**

10    **Cis-N-[1-(5-Chloro-thiophene-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

- [0140]**   R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 5-chloro-thien-2-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

15    **Example 39**

**Cis-N-[1-(1,5-Dimethyl-1H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

- 20    **[0141]**   R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 1,5-dimethyl-1*H*-pyrazol-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.  
The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

25

30

35

40

45

50

55

Example	Name	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>6</sup>	R <sup>7</sup>	R <sup>8</sup>	LC (R.; purity)	MS (ESI)
40	cis-N-[2-Methyl-1-(4-methyl- isothiazole-5-carbonyl)-1,2,3,4- tetrahydro-quinolin-4-yl]-N- phenyl-acetamide	Phenyl	CH <sub>3</sub>	4-methyl-isothiazol-5-yl	CH <sub>3</sub>	H	H	H	H	3.12 min 100%	406 MH+
41	ci-5-[4-(Acetyl-phenyl-amino)-2- methyl-3,4-dihydro-2H- quinoline-1-carbonyl]-thiophene- 2-carboxylic acid dimethylamide	Phenyl	CH <sub>3</sub>	2-carboxylic acid dimethylamide- thiophen-5-yl	CH <sub>3</sub>	H	H	H	H	3.09 min 100%	462 MH+
42	cis-N-[1-(4-Hydroxy-quinoline-6- carbonyl)-2-methyl-1,2,3,4- tetrahydro-quinolin-4-yl]-N- phenyl-acetamide	Phenyl	CH <sub>3</sub>	4-Hydroxy-quinolin-6-yl	CH <sub>3</sub>	H	H	H	H	2.95 min 100%	452 MH+
43	cis-N-[1-(4-tert-Butyl-thiazole-2- carbonyl)-2-methyl-1,2,3,4- tetrahydro-quinolin-4-yl]-N- phenyl-acetamide	Phenyl	CH <sub>3</sub>	4-tert-Butyl-thiazol-2-yl	CH <sub>3</sub>	H	H	H	H	3.54 min 100%	448 MH+
44	cis-N-[1-(2-Ethyl-pyridine-4- carbonyl)-2-methyl-1,2,3,4- tetrahydro-quinolin-4-yl]-N- phenyl-acetamide	Phenyl	CH <sub>3</sub>	2-ethyl-pyridin-4-yl	CH <sub>3</sub>	H	H	H	H	3.04 min 100%	414 MH+

45	cis-N-[1-(3,6-Dichloro-pyridine-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	3,6-Dichloro-pyridin-2-yl	CH <sub>3</sub>	H	H	H	H	3.32 min 100%	454 MH+
46	cis-N-[1-(4-Chloro-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	4-Chloro-2H-pyrazol-3-yl	CH <sub>3</sub>	H	H	H	H	2.94 min 100%	409 MH+
47	cis-2-[4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-isonicotinic acid methyl ester	Phenyl	CH <sub>3</sub>	4-carboxylic acid methyl ester-pyridin-2-yl	CH <sub>3</sub>	H	H	H	H	3.09 min 100%	444 MH+
48	cis-N-[2-Methyl-1-(4-[1,2,4]triazol-4-yl-benzoyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	4-[1,2,4]triazol-4-yl-benzoyl	CH <sub>3</sub>	H	H	H	H	2.95 min 100%	452 MH+
49	cis-N-[1-(2,6-Dimethoxy-pyridine-4-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	2,6-Dimethoxy-pyridin-4-yl	CH <sub>3</sub>	H	H	H	H	3.32 min 100%	446 MH+

50	cis-N-[1-(5-Ethyl-isoxazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	5-Ethyl-isoxazol-3-yl	CH <sub>3</sub>	H	H	H	H	3.14 min 100%	404 MH+
51	cis-N-[2-Methyl-1-(2-tetrazol-1-yl-pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	2-tetrazol-1-yl-pyridin-4-yl	CH <sub>3</sub>	H	H	H	H	3.01 min 100%	454 MH+
52	cis-N-[2-Methyl-1-(5-propyl-isoxazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	5-propyl-isoxazol-3-yl	CH <sub>3</sub>	H	H	H	H	3.27 min 100%	418 MH+
53	cis-N-[1-(5-Isobutyl-2-methyl-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	5-Isobutyl-2-methyl-2H-pyrazol-3-yl	CH <sub>3</sub>	H	H	H	H	3.44 min 100%	445 MH+
54	cis-N-[1-(5-Bromo-furan-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	5-Bromo-furan-2-yl	CH <sub>3</sub>	H	H	H	H	3.27 min 100%	455 MH+

55	cis-N-[2-Methyl-1-(6-phenyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	6-phenyl-pyridin-3-yl	CH <sub>3</sub>	H	H	H	H	3.39 min 100%	462 MH+
56	cis-N-[2-Methyl-1-(2-phenyl-pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	2-phenyl-pyridin-4-yl	CH <sub>3</sub>	H	H	H	H	3.36 min 100%	462 MH+
57	cis-N-[2-Methyl-1-(quinoline-6-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	quinolin-6-yl	CH <sub>3</sub>	H	H	H	H	3.20 min 100%	436 MH+
58	cis-N-[1-(3,4-Dimethoxy-furan-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	3,4-Dimethoxy-furan-2-yl	CH <sub>3</sub>	H	H	H	H	3.19 min 100%	435 MH+
59	cis-N-[2-Methyl-1-(3-methyl-furan-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	3-methyl-furan-2-yl	CH <sub>3</sub>	H	H	H	H	3.19 min 100%	389 MH+

60	cis-N-[1-(2,5-Dimethyl-furan-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	2,5-Dimethyl-furan-3-yl	CH <sub>3</sub>	H	H	H	H	3.36 min 100%	403 MH+
61	cis-N-[1-(2,4-Dimethyl-oxazole-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	2,4-Dimethyl-oxazol-5-yl	CH <sub>3</sub>	H	H	H	H	3.10 min 100%	404 MH+
62	cis-N-[1-(5-Methoxymethyl-furan-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	5-Methoxymethyl-furan-2-yl	CH <sub>3</sub>	H	H	H	H	3.13 min 100%	419 MH+
63	cis-N-[1-(5-Fluoro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	5-Fluoro-pyridin-3-yl	CH <sub>3</sub>	H	H	H	H	3.07 min 100%	404 MH+
64	cis-N-[2-Methyl-1-(quinoline-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	quinoline-2-yl	CH <sub>3</sub>	H	H	H	H	3.22 min 100%	436 MH+

65	cis-N-[2-Methyl-1-(6-methyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	6-methyl-pyridin-3-yl	CH <sub>3</sub>	H	H	H	H	3.00 min 100%	400 MH+
66	cis-N-[2-Methyl-1-(quinoline-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	quinolin-3-yl	CH <sub>3</sub>	H	H	H	H	3.27 min 100%	436 MH+
67	cis-N-[2-Methyl-1-(1H-pyrazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	1H-pyrazol-3-yl	CH <sub>3</sub>	H	H	H	H	2.88 min 100%	375 MH+
68	cis-N-[2-Methyl-1-(2H-pyrazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	2H-pyrazol-3-yl	CH <sub>3</sub>	H	H	H	H	2.88 min 100%	375 MH+
69	cis-N-[1-(5-Isobutyl-isoxazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	5-Isobutyl-isoxazol-3-yl	CH <sub>3</sub>	H	H	H	H	3.38 min 100%	432 MH+



70	cis-N-[2-Methyl-1-(quinoline-4-carbonyl)-1,2,3,4-tetrahydroquinolin-4-yl]-N-phenylacetamide	Phenyl	CH <sub>3</sub>	quinolin-4-yl	CH <sub>3</sub>	H	H	H	H	3.20 min 100%	435 MH+
71	cis-N-[2-Methyl-1-(6-methylpyridine-2-carbonyl)-1,2,3,4-tetrahydroquinolin-4-yl]-N-phenylacetamide	Phenyl	CH <sub>3</sub>	6-methyl-pyridin-2-yl	CH <sub>3</sub>	H	H	H	H	3.06 min 100%	400 MH+
72	cis-N-[2-Methyl-1-(quinoxaline-5-carbonyl)-1,2,3,4-tetrahydroquinolin-4-yl]-N-phenylacetamide	Phenyl	CH <sub>3</sub>	quinoxalin-5-yl	CH <sub>3</sub>	H	H	H	H	3.15 min 100%	437 MH+
73	cis-N-[1-(3-Methoxy-thiophene-2-carbonyl)-2-methyl-1,2,3,4-tetrahydroquinolin-4-yl]-N-phenylacetamide	Phenyl	CH <sub>3</sub>	3-Methoxy-thiophen-2-yl	CH <sub>3</sub>	H	H	H	H	3.25 min 100%	421 MH+
74	cis-N-[1-(5-tert-Butyl-2-methylfuran-3-carbonyl)-2-methyl-1,2,3,4-tetrahydroquinolin-4-yl]-N-phenylacetamide	Phenyl	CH <sub>3</sub>	5-tert-Butyl-2-methyl-furan-3-yl	CH <sub>3</sub>	H	H	H	H	3.65 min 100%	445 MH+

75	cis-N-[1-(5-Ethyl-2-methyl-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	5-Ethyl-2-methyl-2H-pyrazol-3-yl	CH <sub>3</sub>	H	H	H	H	3.23 min 100%	417 MH+
76	cis-N-[2-Methyl-1-([1,2,5]thiadiazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	[1,2,5]thiadiazol-3-yl	CH <sub>3</sub>	H	H	H	H	3.03 min 100%	392 MH+
77	cis-N-[2-Methyl-1-(2-methyl-5-propyl-2H-pyrazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide	Phenyl	CH <sub>3</sub>	2-methyl-5-propyl-2H-pyrazol-3-yl	CH <sub>3</sub>	H	H	H	H	3.35 min 100%	431 MH+

**Example 78****Cis-N-[2-Methyl-1-(pyridine-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0142] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: pyridin-2-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H

The titled compound was prepared according to protocol A, using intermediate A and appropriate reactants.

<sup>1</sup>H NMR [(CDCl<sub>3</sub>): 8.5 (s, 1H), 7.4-7.1 (m, 7H), 6.95 (d, 1H), 6.8 (m, 1H), 6.5 (d, 1H), 5.55 (m, 1H), 4.8 (m, 1H), 2.3 (m, 1H), 2.0 (s, 3H), 1.25 (m, 1H), 1.1 (d, 3H).

HPLC: r<sub>T</sub>: 8.6 min.

MS positive ESI: m/z (m+H)<sup>+</sup>= 386

**Example 79****Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-benzyl-acetamide**

[0143] R<sup>1</sup>: benzyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H

The titled compound was prepared according to protocol A, using intermediate Cis-B and appropriate reactants.

<sup>1</sup>H NMR [(CDCl<sub>3</sub>): 7.4 (m, 1H), 7.2 (m, 8H), 7.0 (m, 1H), 6.9 (m, 1H), 6.5 (m, 1H), 5.6 (d, 0.5H), 5.0 (m, 1H), 4.8 (m, 1H), 3.9 (m, 0.5H), 2.5 (m, 1H), 2.3, 2.4 (s, 3H), 1.5 (m, 1H), 1.2 (m, 3H)

HPLC: r<sub>T</sub>: 3.17 min.

MS positive ESI: m/z (m+H)<sup>+</sup>= 399

**Example 80****Cis-N-Benzyl-N-[2-methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide**

[0144] R<sup>1</sup>: benzyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: thien-2-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate Cis-B and appropriate reactants.

<sup>1</sup>H NMR [(CDCl<sub>3</sub>): 7.5-6.7 (m, 12H), 5.6 (d, 0.5H), 4.9 (m, 1H), 4.7 (m, 1H), 3.9 (d, 0.5H), 2.5 (m, 1H), 2.31, 2.29 (s, 3H), 1.5 (m, 1H), 1.2 (app.d, 3H)

HPLC: r<sub>T</sub>: 3.07 min.

MS positive ESI: m/z (m+H)<sup>+</sup>= 405

**Example 81****Trans-N-Benzyl-N-[2-methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide**

[0145] R<sup>1</sup>: benzyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: thien-2-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using intermediate Trans-B and appropriate reactants.

<sup>1</sup>H NMR [(CDCl<sub>3</sub>): 7.5-6.9 (m, 12H), 6.2 (t, 1H), 5.0 (m, 1H), 4.6 (d, 1H), 4.4 (d, 1H), 2.2 (s, 3H), 2.1 (m, 2H), 1.3 (d, 3H)

HPLC: r<sub>T</sub>: 3.03 min.

MS positive ESI: m/z (m+H)<sup>+</sup>= 405

**Example 82****Cis-N-Cyclohexyl-N-[2-methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide**

[0146] R<sup>1</sup>: cyclohexyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: thien-2-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using appropriate reactants.

<sup>1</sup>H NMR [(CDCl<sub>3</sub>): 7.5-6.7 (m, 7H), 5.2 (m, 1H), 3.7 (m, 1H), 2.8 (m, 1H), 2.2 (s, 3H), 2.2-1.2 (m, 12 H), 1.3 (d, 3H)

HPLC: r<sub>T</sub>: 3.27 min.

MS positive ESI: m/z (m+H)<sup>+</sup>= 398

**Example 83****Cis-N-(1-Benzoyl-6-methoxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide**

[0147] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: O-CH<sub>3</sub>, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol A, using appropriate reactants and starting from 4-chloro-6-methoxy-quinoline.

(0.045g, 36% yield)

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ 7.5 (m, 2H), 7.4 (m, 3H), 7.3 (m, 1H), 7.2 (m, 2H), 7.1 (m, 2H), 6.9 (m, 1H), 6.5 (m, 1H), 6.4 (m, 1H), 5.4 (m, 1H), 4.6 (m, 1H), 3.6 (s, 3H), 2.6 (m, 1H), 1.9 (s, 3H), 1.2 (m, 1H), 1.0 (d, 3H).

HPLC: *r*<sub>T</sub>: 10.05, 98.8% purity

MS positive ESI: *m/z* (*m*+*H*)<sup>+</sup> = 415

#### Example 84

##### Cis-N-(1-Benzoyl-6-hydroxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide

[0148] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: OH, R<sup>7</sup>: H, R<sup>8</sup>: H.

To a solution of 79 mg (0.19mM) of Cis-N-(1-Benzoyl-6-methoxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide in 3ml of dichloromethane were dropwise added 50 µl of boron tribromide. After 3 hours, ice was added and the mixture was extracted by ethyl acetate. The organic layer was dried by sodium sulfate and evaporated under reduced pressure to give a white solid that was purified by silicagel with dichloromethane/methanol 95/5 as eluent. (0.073g, 96% yield).

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ +9.4 (s, 1H) 7.5 (m, 2H), 7.4 (m, 3H), 7.3 (m, 1H), 7.2 (m, 2H), 7.1 (m, 2H), 6.8 (s, 1H), 6.3 (s, 2H), 5.4 (m, 1H), 4.6 (m, 1H), 2.4 (m, 1H), 1.9 (s, 3H), 1.3 (m, 1H), 1.0 (m, 3H).

HPLC: *r*<sub>T</sub>: 8.58, 100% purity

MS positive ESI: *m/z* (*m*+*H*)<sup>+</sup> = 401

#### Example 85

##### Cis-N-(1-Benzoyl-6-chloro-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide

[0149] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: Cl, R<sup>7</sup>: H, R<sup>8</sup>: H.

To a solution of 4,6-dichloroquinoline (4g) in methanol (50ml) was added 1.1 eq (1.1g) of sodium methoxyde. The mixture was heated under reflux for 17 hours. After cooling, the reaction mixture was evaporated under reduce pressure, dissolved in dichloromethane (80ml) and washed with water (80ml). The aqueous layer was extracted by 80ml of dichloromethane. The organic layers were combined and dried by sodium sulfate. The solvent was evaporated to give a brown oil which was chromatographed in silicagel using CH<sub>2</sub>Cl<sub>2</sub> and CH<sub>2</sub>Cl<sub>2</sub>/MeOH 99/1 and then 98/2. The fractions containing 4-methoxy-6-chloro-quinoline (**3a**) were combined and evaporated to give a brown solid (2.5 g; yield = 63%).

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ 8.8 (m, 1H), 8.10 (m, 1H), 7.98 (m, 1H), 7.75 (m, 1H), 7.10 (m, 1H), 4.05 (s, 1H).

MS positive ESI: *m/z* (*m*+*H*)<sup>+</sup> = 193 (75%), 195 (25%)

HPLC: *r*<sub>T</sub>: 4.97, 100% purity

[0150] To a solution of 4-methoxy-6-chloro-quinoline (2.5g) in tetrahydrofuran (60 ml), under nitrogen and under cooling at -78°C (in an acetone-dry ice bath) and stirring, a solution of methyl magnesium chloride (3 mol/L) in tetrahydrofuran was dropwise added (21.4ml) for 1 hour and then benzoyl chloride (7.5 ml) was dropwise added for 30 minutes. The mixture was stirred for 2 hours. Methanol (130 ml) and then chlorhydric acid 1 mol/L (130 ml) were added and the reaction mixture was stirred overnight. Tetrahydrofuran and methanol were evaporated under reduce pressure and the aqueous layer was extracted by dichloromethane (200 ml). After separation, the organic layer was dried by sodium sulfate, evaporated under reduced pressure and chromatographed in silicagel with dichloromethane as eluent.

The fractions containing 1-benzoyl-6-chloro-2-methyl-2,3-dihydro-1*H*-quinolin-4-one (**3b**) were combined and evaporated under reduced pressure to give a yellow solid (0.7g ; yield =18%).

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ 8-7 (m, 9H), 5(m, 1H), 3.33 (m, 1H), 2.60 (m, 1H), 1.20 (m, 1H).

HPLC: *r*<sub>T</sub>: 10.50, 84% purity

MS positive ESI: *m/z* (*m*+*H*)<sup>+</sup> = 299 (75%), 301 (25%)

[0151] To a solution of 1-benzoyl-6-chloro-2-methyl-2,3-dihydro-1*H*-quinolin-4-one (0.7g), aniline (3eq) and triethylamine (5eq) in dichloroethane (30 ml), cooled at 0°C in an ice bath, 1 eq of a solution of titanium chloride in toluene (1 mol/l) was dropwise added for 15 min. The mixture was heated at 85°C during 4H30. The mixture was washed by an aqueous solution of sodium bicarbonate. The organic layer was separated, dried over sodium sulfate and the solvent was removed under reduce pressure to give a brown oil (1g) (6-chloro-2-methyl-4-phenylimino-3,4-dihydro-2*H*-quinolin-1-yl)-phenyl-methanone (**3c**), which was not purified.

[0152] To a solution of 6-chloro-2-methyl-4-phenylimino-3,4-dihydro-2*H*-quinolin-1-yl)-phenyl-methanone (0.87 g) in glacial acetic acid (35 ml) was added in small amount sodium triacetoxymethylborohydride (6eq) under stirring at room temperature. The reaction was stirred during 2 hours.

Ice (100g) was added and the mixture was basified by an aqueous solution of sodium hydroxyde until pH 6 (1 mol/l, 200ml). The aqueous solution was extracted by dichloromethane (400ml). The organic layer was separated, dried over sodium sulfate and the solvent was removed under reduced pressure to give an amber solid. A mixture of cis/trans (6-chloro-2-methyl-4-phenylamino-3,4-dihydro-2H-quinolyn-1-yl)-phenyl-methanone (86/14) was obtained after a first chromatography on silica gel (dichloromethane).

Cis-(6-chloro-2-methyl-4-phenylamino-3,4-dihydro-2H-quinolyn-1-yl)-phenyl-methanone was obtained after a second chromatography on silica gel (cyclohexane /ethylacetate 95/5) (90 mg, yield = 10.3%).

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ 7.3 (m, 5H), 7.12 (m, 3H), 6.99 (m, 1H), 6.78(m,2H), 6.62(m,1H), 6.51 (m, 1H), 6.04 (m, NH), 4.78 (m, 1H), 4.58 (m, 1H), 2.72 (m,1H), 1.18 (m,3H).

HPLC: r<sub>T</sub>: 11.70, 97% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 377

**[0153]** A solution of Cis-(6-chloro-2-methyl-4-phenylamino-3,4-dihydro-2H-quinolyn-1-yl)-phenyl-methanone (40mg) in dimethylformamide (1 ml) and diisopropylethylamine (3eq) was stirred during 8 hours. Acetylchloride (6 eq) was added and the reaction mixture was stirred for 72H. The solvent was evaporated in reduced pressure and the residue was dissolved in dichloromethane (10ml). The organic layer was washed by water (10ml), dried over sodium sulfate and evaporated under reduced pressure. The crude material was purified by chromatography on silica gel (dichloromethane/methanol 100/0 and 99/1) to afford the title compound (0.025g, 57% yield).

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ 7.5 (m, 2H), 7.4 (m, 5H), 7.3 (m, 2H), 7.2 (m, 2H), 7.0 (m, 1H), 6.5 (m, 1H), 5.4 (m, 1H), 4.6 (m, 1H), 2.7 (m, 1H), 1.9 (s, 3H), 1.2 (m, 1H), 1.0 (m, 3H).

HPLC: r<sub>T</sub>: 10.9, 99.2% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 419 (75%), 421 (25%)

#### Example 86

##### Cis-N-(1-Benzoyl-7-chloro-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide

**[0154]** R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: Cl, R<sup>8</sup>: H.

The titled compound was prepared according to protocol B, using appropriate reactants and starting from 4-methoxy-7-chloro-quinoline.

(1.04g, 70% yield)

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ 7.5 (m, 4H), 7.4 (m, 3H), 7.3 (m, 2H), 7.2 (m, 3H), 6.6 (m, 1H), 5.4 (m, 1H), 4.6 (m, 1H), 2.7 (m, 1H), 1.9 (s, 3H), 1.3 (m, 1H), 1.0 (m,3H).

HPLC: r<sub>T</sub>: 10.9, 95% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 419 (75%), 421 (25%)

#### Example 87

##### N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-prop-2-ynyl-acetamide

**[0155]** R<sup>1</sup>: propynyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

To a solution of 1-benzoyl-2-methyl-2,3-dihydro-1H-quinolin-4-one (corresponds to a compound of general formula (3b) wherein R<sup>3</sup> is phenyl, R<sup>4</sup> is CH<sub>3</sub> and R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> are H, prepared as disclosed in example 46, according to protocol B, starting from 4-chloro-quinoline, (0.3g), triethylamine (0.92g), propargyl amine (0.125g) in dichloroethane (20 ml), cooled at -5°C in an ice/sodium chloride bath, a solution of titanium chloride in toluene 1 mol/l was dropwise added for 10 minutes. The mixture was stirred over night at room temperature and then washed with dichloromethane and an aqueous solution of potassium carbonate. The organic layer was separated, dried over sodium sulfate and the solvent was removed under reduced pressure to give (2-methyl-4-prop-2-ynylimino-3,4-dihydro-2H-quinolin-1-yl)-phenyl-methanone (brown oil) (0.5g) which was not purified, but directly used in the next step

MS positive ESI: m/z (m+H)<sup>+</sup>= 303

**[0156]** To a solution of (2-methyl-4-prop-2-ynylimino-3,4-dihydro-2H-quinolin-1-yl)-phenyl-methanone (0.5g) in acetic acid (20 ml), sodium triacetoxyborohydride was added in small amounts (1.75g) under stirring at room temperature. The reaction mixture was then stirred at room temperature for one hour and poured on 100 ml of cold water under stirring. The desired compound was filtered after precipitation. The compound was dissolved in ethyl acetate and washed with water. The organic layer was separated and dried over sodium sulfate. The solvent was removed under reduced pressure to give a brown oil which was purified by flash chromatography with cyclohexane/ethyl acetate (95/5) as eluent and then cyclohexane/ethyl acetate (90/10) to give a mixture of diastereoisomers cis-trans (50-50) of (2-methyl-4-prop-2-ynylamino-3,4-dihydro-2H-quinolin-1-yl)-phenyl-methanone (0.06g, 17.6% yield).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 6.95-7.4 (m, 7H), 6.9 (t, 1H), 6.5 (d, 1H), 4.85 (m, 1H), 4.15 (m, 0.5H), 3.9 (m, 0.5H), 3.7 (s, 1H),

3.5 (d, 0.5), 3.35 (d, 0.5H), 2.8 (m, 0.5), 2.5 (m, 0.5H), 2.3 (m, 1H), 1.75 (m, 0.5H), 1.1-1.4 (m, 3.5H).

HPLC: Cis  $r_T$ : 5.86 min, 46.96% , purity

Trans  $r_T$ : 6.27 min, 47.12%, purity

MS positive ESI:  $m/z$  ( $m+H$ ) $^+$ = 305

5 [0157] To a solution of (2-methyl-4-prop-2-ynylamino-3,4-dihydro-2H-quinolin-1-yl)-phenyl-methanone (0.06 g), di-isopropyl ethyl amine (0.038 g) in dioxane ( 5 ml), anhydride acetic (0.2g) was added under stirring. The mixture was heated under reflux over the night. After cooling dichloromethane and an aqueous solution of potassium carbonate were added. The organic layer was separated, and then dried over sodium sulfate. Organic solvent was removed under reduce pressure to give the crude compound (0.15 g). Purification by chromatography was performed with dichloromethane/methanol (99/1) as eluent to give the desired title compound (0.04g, 58.8% yield).

MS positive ESI:  $m/z$  ( $m+H$ ) $^+$ = 347

HPLC: Cis  $r_T$ : 7.628 min, 49.05% , purity

Trans  $r_T$ : 7.915 min, 50.95%, purity

#### 15 Example 88

##### **Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-(4-methoxy-phenyl)-acetamide**

[0158] R<sup>1</sup>: 4-methoxy-phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

20 The titled compound was prepared according to protocol B, as disclosed in example 48, using 4-methoxy-phenylamine instead of propargyl-amine in the first step.

(0.07g, 63.6% yield)

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]:  $\delta$  7.45 (m, 1H), 7.1-7.4 (m, 9H), 7 (m, 1H), 6.9 (m, 1H), 6.5 (m, 1H), 5.5 (m, 1H), 64.65 (m, 1H), 3.8 (s, 3H), 2.7 (m, 1H), 1.9 (s, 3H), 1.25 (m, 1H), 1.05 (m, 3H).

25 HPLC:  $r_T$ : 10.003, 93.8% purity

MS positive ESI:  $m/z$  ( $m+H$ ) $^+$ = 415

#### Example 89

##### 30 **Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-(4-hydroxy-phenyl)-acetamide**

[0159] R<sup>1</sup>: 4-hydroxy-phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

[0160] To a solution of Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-(4-methoxy-phenyl)-acetamide (0.1g) in dichloromethane (5ml), boron tribromide (0.18g) was added under stirring and under cooling at -78°C in an acetone-dry ice bath.

35 The mixture was stirred over night at room temperature, and then washed with dichloromethane (20ml) and water (20ml). The organic layer was separated, dried over sodium sulfate and the solvent was removed under reduce pressure. The crude compound was purified by flash chromatography on silica gel with an eluent dichloromethane/methanol (97/3) to give a solid (0.045g, 46.6% yield).

40 <sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]:  $\delta$  9.7 (s, 1H), 7.1-7.4 (m, 9H), 6.9 (m, 1H), 6.8 (m, 2H), 6.5 (m, 1H), 5.45 (m, 1H), 4.65 (m, 1H), 2.55 (m, 1H), 1.9 (s, 3H), 1.25 (m, 1H), 1.05 (m, 3H).

HPLC:  $r_T$ : 8.57, 100% purity

MS positive ESI:  $m/z$  ( $m+H$ ) $^+$ = 401

#### 45 Example 90

##### **Cis-[4-[Acetyl-(1-benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-amino]-phenyl]-acetic acid ethyl ester**

[0161] R<sup>1</sup>: 4-methyl acid ethyl ester-phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

50 The titled compound was prepared according to protocol B, as disclosed in example 48, using 4-methyl acid ethyl ester-phenylamine.

(0.1g, 30.4% yield)

<sup>1</sup>H NMR [CDCl<sub>3</sub>]:  $\delta$  7.1-7.4 (m, 11H), 6.9 (m, 1H), 6.5 (m, 1H), 6.65 (m, 1H), 4.8 (m, 1H), 4.15 (m, 2H), 3.65 (s, 2H), 2.35 (m, 1H), 2.05 (m, 3H), 1.25 (m, 4H), 1.1 (m, 3H).

55 HPLC:  $r_T$ : 10.42, 95.73% purity

MS positive ESI:  $m/z$  ( $m+H$ ) $^+$ = 471

**Example 91****Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-malonamic acid methyl ester**

[0162] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>2</sub>-COOCH<sub>3</sub>, R<sup>3</sup>: phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol B, using appropriate reactants. (0.1g, 77.5% yield)

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ 7.65 (m, 1H), 7.05-7.6 (m, 11H), 6.95 (m, 1H), 6.5 (m, 1H), 5.5 (m, 1H), 4.65 (m, 1H), 3.6 (s, 3H), 3.35 (m, 2H), 2.55 (m, 1H), 1.2 (m, 1H), 1 (m, 3H).

HPLC: r<sub>T</sub>: 10.3, 100% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 442

**Example 92****Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-malonamic acid**

[0163] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>2</sub>-COOH, R<sup>3</sup>: phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

To a solution of Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-malonamic acid methyl ester (0.031g) in a mixture of methanol (1ml) and water (1ml), 0.0067g of lithium hydroxide was added under stirring at room temperature, over night. In order to complete the reaction, 0.0067g of lithium hydroxide was added and the mixture was stirred over night. Then dichloromethane and an aqueous solution of sodium bicarbonate were added, the organic layer was separated, dried over sodium sulfate, the solvent was removed under reduce pressure to give a solid which was purified by flash chromatography with dichloromethane/methanol (95/5) as eluent to give a solid (0.01g, 33.3% yield).

HPLC: r<sub>T</sub>: 9.09, 100% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 427

**Example 93****Cis-N-[2-Methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide**

[0164] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: pyridin-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

[0165] A mixture of 4-chloroquinoline (10g) and aniline (11.4g) was warmed to 150°C for 30 min. A solution of sodium hydroxide 1 N (200ml) was added. After extraction with ethyl acetate (1L), the organic layer was separated and dried over sodium sulfate. The solvent was removed to afford an amorphous material, which was crystallized in cyclohexane. Phenyl-quinolin-4-yl-amine (13.7g) was obtained after filtration.

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ 8.95 (s, 1H), 8.45 (d, 1H), 8.35 (d, 1H), 7.9 (m, 1H), 7.7 (m, 1H), 7.55 (m, 1H), 7.4 (m, 4H), 7.15 (m, 1H), 6.95 (m, 1H).

[0166] To a solution of phenyl-quinolin-4-yl-amine (13g) and N-ethyl-N,N-diisopropylamine (1.5 eq) in dioxane (105 ml) was added acetic anhydride (28.1 ml, 5 eq). The solution was warmed to reflux of dioxane for 5 days. After cooling, dichloromethane was added and the solution was washed by water and sodium hydroxide 1N. The organic layer was separated and dried over sodium sulfate. The solvent was removed and the mixture was purified by chromatography on silica gel (dichloromethane/ ethanol (95:5)) to afford N-phenyl-N-quinolin-4-yl-acetamide (10g).

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ 9.0 (m, 1H), 8.1 (m, 2H), 7.8 (m, 1H), 7.65 (m, 1H), 7.55-7.3 (m, 6H), 2.2-2.0 (m, 3H).

[0167] A solution of N-phenyl-N-quinolin-4-yl-acetamide (2g) and benzyle bromide (1.3g) in acetone was warmed to reflux for 5 days. The solvent was removed to afford an amorphous material, which was crystallized in diethyl ether. 4-[Acetyl-(phenyl)-amino]-1-benzylquinolinium bromide (2g) was obtained by filtration.

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ 9.75 (d, 1H), 8.45 (q, 2H), 8.2 (m, 2H), 8.0 (m, 1H), 7.65 (m, 2H), 7.55-7.35 (m, 8H), 6.3 (s, 2H), 2.2 (s, 3H).

[0168] To a suspension of 4-[(acetyl)(phenyl)amino]-1-benzylquinolinium bromide (200mg) in tetrahydrofuran, under cooling at -78°C in an acetone-dry ice bath and under stirring, a solution of methyl magnesium chloride at 3M/1 was dropwise added. The suspension became homogenous. At -78°C, a mixture of methanol and water were added, and then extracted with ethyl acetate. The organic layer was separated, dried over sodium sulfate. The solvent was removed under reduced pressure to give the desired compound (150mg).

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ 7.5-7.15 (m, 10H), 7.0-6.9 (m, 2H), 6.5 (m, 2H), 6.1 (m, 1H), 4.3 (m, 1H), 2.1 (2m, 3H), 1.15 (d, 3H).

[0169] A solution of N-(1-benzyl-2-methyl-1,2-dihydroquinolin-4-yl)-N-phenylacetamide (100mg) and (R,R)-(-)-1,2-bis[(o-methoxyphenyl)(phenyl)phosphino]ethane(1,5-cyclooctadiene)rhodium (I) tetrafluoroborate (20mg) in methanol (6ml) was stirred under 20 bars of hydrogen overnight. Dichloromethane was added and the organic layer was

washed by water and dried over sodium sulfate. The solvent was removed and the mixture was purified by chromatography on silica gel (dichloromethane / ethanol (99:1)) to afford N-(1-benzyl-2-methyl-1,2,3,4-tetrahydroquinolin-4-yl)-N-phenylacetamide (80mg).

<sup>1</sup>H NMR [(CDCl<sub>3</sub>): δ 7.35-7.1 (m, 11H), 7.0 (t, 1H), 6.7 (t, 1H), 6.4 (d, 1H), 6.2 (m, 1H), 4.4 (dd, 2H), 3.6 (m, 2H), 1.9 (m+s, 4H), 1.5 (m, 1H), 1.05 (d, 3H).

**[0170]** A solution of N-(1-benzyl-2-methyl-1,2,3,4-tetrahydroquinolin-4-yl)-N-phenylacetamide (50mg) and ammonium formate (85mg) and palladium on activated carbon 10% in ethanol (2ml) were warmed to reflux for 1 hour. The solution was filtered on celite and the solvent was removed to afford N-(2-methyl-1,2,3,4-tetrahydroquinolin-4-yl)-N-phenylacetamide (40 mg).

<sup>1</sup>H NMR [(CDCl<sub>3</sub>): δ 7.25 (m, 4H), 7.0 (m, 3H), 6.75 (t, 1H), 6.45 (d, 1H), 6.3 (m, 1H), 3.55 (m, 2H), 1.9 (m+s, 4H), 1.35 (m, 1H), 1.1 (d, 3H).

**[0171]** Nicotinoyl chloride (1.1 eq) was added to a solution of N-(2-methyl-1,2,3,4-tetrahydroquinolin-4-yl)-N-phenylacetamide (100 mg) and N-ethyl-N,N-diisopropylamine (2.2 eq) in dioxane (1 ml). The mixture was stirred overnight at room temperature. Dichloromethane was added and the organic layer was washed by water and sodium carbonate solution and dried over sodium sulfate. The solvent was removed and the mixture was purified by chromatography on silica gel (dichloromethane and ethanol (99:1)) to afford the title compound (30 mg).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 8.6 (s, 1H), 8.5 (d, 1H), 7.4-7.2 (m, 8H), 7.1 (m, 1H), 6.9 (t, 1H), 6.45 (d, 1H), 5.6 (m, 1H), 4.8 (m, 1H), 2.3 (m, 1H), 2.0 (s, 3H), 1.1 (d+m, 4H).

HPLC:  $r_f$ : 8.11, 97.2% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 386

#### Example 94

##### Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-Cyclopropylacetamide

**[0172]** R<sup>1</sup>: cyclopropyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: phenyl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol C, using appropriate reactants.

(0.24g, 47% yield)

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ 6.75-7.4 (m, 8H), 6.5 (m, 1H), 5.35 (m, 1H), 4.7 (m, 1H), 2.8 (m, 1H), 2.5 (m, 1H), 2.3 (m, 3H), 1.9 (m, 1H), 1.3 (m, 3H), 1.1 (m, 1H), 1 (m, 1H), 0.75 (m, 1H), 0.55 (m, 1H).

HPLC: 99.55%, purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 349

#### Example 95

##### Cis-N-Cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide

**[0173]** R<sup>1</sup>: cyclopropyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: pyridin-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

**[0174]** A mixture of 4-chloroquinoline (13.9 g) and cyclopropyl amine (9.7 g) was heated at 180 °C under stirring for 16 hours. After cooling, a solution of sodium hydroxide 1N (200 ml) was added. After extraction with dichloromethane (200 ml), the organic layer was separated and dried over sodium sulfate. The solvent was removed under reduce pressure to afford an oil, which was crystallized in acetonitrile (50ml).

Cyclopropyl-quinolin-4-yl-amine (6.6 g; 42% yield) was obtained after filtration.

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ 8.5 (m, 1H), 8.2 (m, 1H), 7.8 (m, 1H), 7.6 (m, 1H), 7.4 (m, 1H), 6.8 (m, 1H), 2.6 (m, 1H), 0.85 (m, 2H), 0.6 (m, 2H). HPLC:  $r_f$ : 5.7, 100% purity

**[0175]** Acetic anhydride (18.3 g) was added to a solution of cyclopropyl-quinolin-4-yl-amine (6.6g) in pyridine (50 ml) and the mixture was heated under reflux over night. After cooling, the solvent was removed under reduce pressure and then the mixture was washed with dichloromethane (200 ml) and an aqueous solution of sodium carbonate (200 ml). The organic layer was separated and dried over sodium sulfate. The solvent was removed under reduce pressure and the crude compound was purified by flash chromatography on silicagel using dichloromethane/ methanol (98/2) as eluent to give cyclopropyl-quinolin-4-yl-acetamide (8 g, 98.7 % yield).

<sup>1</sup>H NMR [(CD<sub>3</sub>)<sub>2</sub>SO]: δ (T= 400 K) 8.9 (m, 1H), 8.1 (m, 1H), 7.7-7.85 (m, 2H), 7.6 (m, 1H), 7.35 (m, 1H), 3.4 (m, 1H), 2.1 (s, 3H), 0.85 (m, 2H), 0.55 (m, 2H). MS positive ESI: m/z (m+H)<sup>+</sup>= 227

**[0176]** Benzyl bromide (16.71 g) was added to a solution of cyclopropyl-quinolin-4-yl-acetamide (7.37 g) in acetone (50 ml). The mixture was heated 3 days under reflux.

**[0177]** After cooling, the solid was filtered, washed with acetone (20 ml) and dried to give the 4-(acetyl-cyclopropyl-amino)-1-benzyl-quinolinium bromide (10.6 g, 81.7% yield).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 10.7 (d, 1H), 8.4 (m, 1H), 8.2 (m, 1H), 8.05 (m, 1H), 7.95 (m, 1H), 7.8 (m, 1H), 7.3-7.5 (m, 5H), 6.6



(s, 2H), 3.6 (m, 1H), 2.6 (s, 3H), 1.2 (m, 2H), 0.75 (m, 2H).

HPLC:  $r_T$ : 6.69, 100% purity

[0178] To a mixture of 4-(acetyl-cyclopropyl-amino)-1-benzyl-quinolinium bromide (9.6 g) in tetrahydrofuran (200 ml), under nitrogen and under cooling at 0°C, a solution of methyl magnesium chloride 1 Mol/l (12 ml) was added during 15 minutes. After 2 hours, a solution of ammonium chloride in methanol (100 ml) was added. The solvent was removed under reduce pressure and then washed with dichloromethane and water. The organic layer was separated and the solvent was removed under reduce pressure. The crude compound was purified by flash chromatography on silicagel with an eluent of dichloromethane/ methanol (98/2) to give the N-(1-benzyl-2-methyl-1,2-dihydro-quinolin-4-yl)-N-cyclopropyl-acetamide (7.9g, 98.6% yield).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.3 (m, 6H), 7.1 (m, 1H), 6.95 (m, 1H), 6.65 (m, 1H), 6.5 (m, 1H), 5.45 (m, 1H), 4.6 (m, 1H), 4.35 (m, 1H), 4.2 (m, 1H), 3.2 (m, 1H), 2.05 (m, 3H), 1.2 (m, 3H), 0.8 (m, 2H), 0.55 (m, 2H).

HPLC:  $r_T$ : 10.9, 100% purity

[0179] To a solution of N-(1-benzyl-2-methyl-1,2-dihydro-quinolin-4-yl)-N-cyclopropyl-acetamide (7.5 g) in methanol (160 ml) and tetrahydrofuran (375 ml), nickel chloride hexahydrate (5.4 g) was added. After cooling at 0°C, sodium borohydride (3.41 g) was added in small portions, and then the mixture was stirred at room temperature overnight. Dichloromethane and water were added to the mixture. The organic layer was separated and the solvent was removed under reduce pressure. The crude compound was purified by flash chromatography on silicagel using cyclohexane/ methanol (90/10), then (80/20), then (70/30) to give the Cis-N-(1-benzyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-cyclopropyl-acetamide (3.9 g, 51.7% yield).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.3 (m, 5H), 6.95 (m, 1H), 6.7 (m, 1H), 6.55 (m, 1H), 6.45 (m, 1H), 5.9 (m, 1H), 4.5 (dd, 2H), 3.65 (m, 1H), 2.65 (m, 1H), 2.5 (m, 1H), 2.4 (s, 3H), 1.25 (m, 3H), 0.6-0.9 (m, 4H).

[0180] To a solution of Cis-N-(1-benzyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-cyclopropyl-acetamide (3.8 g) in ethanol (150 ml), ammonium formate (7.2 g) and few milligrams of palladium on charcoal (10 % Pd/C) were added. The mixture was heated under reflux for 2 hours. After cooling the mixture was filtered on celite and the solvent was removed under reduce pressure to give Cis-N-cyclopropyl-N-(2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-acetamide (2.56 g, 92.4% yield) which was not purified.

HPLC:  $r_T$ : 7.62, 99.06% purity

[0181] Nicotinoyl chloride hydrochloride (1.46 g) was added to a solution of Cis-N-cyclopropyl-N-(2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-acetamide (0.4 g) in pyridine (20 ml). The mixture was heated at 50°C for 3 hours. After cooling, an aqueous solution of sodium carbonate was added and the mixture was extracted with ethyl acetate. The organic layer was dried over sodium sulfate, and then the solvent was removed under reduce pressure. The crude compound was crystallized in diethyl ether to give Cis-N-cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide (0.3 g, 52.6 % yield).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 8.5 (s, 1H), 8.4 (3, 1H), 7.15-6.7 (m, 5H), 6.4 (d, 1H), 5.4 (m, 1H), 4.8 (m, 1H), 2.6 (m, 2H), 2.3 (s, 3H), 1.9 (m, 1H), 1.2 (m, 3H), 1.0-0.7 (m, 4H) HPLC:  $r_T$ : 6.39, 98.3% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 350

#### Example 96 and 97

(+)-Cis-N-cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide (example 57) and (-)-Cis-N-cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide (example 58)

[0182] The racemic Cis-N-cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide (0.05 g) was separated by preparative high performance liquid chromatography following the conditions:

Column: Chiralpak AS-H 20\*250 mm, 5μ  
Mobile phase: 70 Hexane (0.2% Diethyl amine)  
30 EtOH (0.2% Diethyl amine)  
Detection: 254 nm

After separation, and concentration, the following products were obtained:

(+)-Cis-N-cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide (0.147 g)

HPLC  $r_T$ : 9.16 min, ee> 99 %, purity

[α]<sub>D</sub><sup>20</sup>= + 249°, (c=0.122, MeOH)

(-)-Cis-N-cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide (0.147 g)

HPLC  $r_T$ : 9.18 min, ee= 94.26 %, purity

$[\alpha]_D^{20} = -227^\circ$ , (c=0.079, MeOH)

### Example 98

#### Cis-N-Cyclopropyl-N-[2-methyl-1-(3-methyl-isoxazole-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide

[0183] R<sup>1</sup>: cyclopropyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: 3-methyl-isoxazol-5-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol C, using appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.15-6.7 (m, 5H), 5.35 (m, 1H), 4.9 (m, 1H), 2.8 (m, 1H), 2.5 (m, 1H), 2.4-2.2 (m, 6H), 1.9 (m, 1H), 1.3 (m, 3H), 1.0-0.7 (m, 4H).

HPLC:  $r_T$ : 7.97, 98.6% purity

MS positive ESI: m/z (m+H)<sup>+</sup> = 354

### Example 99

#### Cis-N-Phenyl-N-[1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide

[0184] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: thien-2-yl, R<sup>4</sup>: H, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

[0185] To a solution of N-phenyl-N-quinolin-4-ylacetamide (2g) and Nickel dichloride (7.6 mmol) in methanol, cooled at 0°C, was added sodium borohydride (20 eq). The solution was then stirred overnight, at room temperature. The solvent was removed to give an amorphous material, which is crystallized in diethyl ether. After filtration, dichloromethane was added and the solution was washed with water. The organic layer was separated and dried over sodium sulfate. The solvent was removed to obtain an amorphous material. The product was purified by chromatography on silica gel using a mixture of cyclohexane and ethyl acetate (95:5) as eluent to give a N-phenyl-N-1,2,3,4-tetrahydro-quinolin-4-ylacetamide (380 mg).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.45 (m, 2H), 7.2 (m, 2H), 7.0 (m, 1H), 6.9 (m, 1H), 6.8 (s, 1H), 6.6 (t, 1H), 6.3 (d, 1H), 6.1 (t, 1H), 3.55 (s, 1H), 3.15 (m, 1H), 2.75 (m, 1H), 2.05 (s, 3H), 1.95 (m, 1H), 1.75 (m, 1H).

[0186] To a solution of N-phenyl-N-1,2,3,4-tetrahydroquinolin-4-ylacetamide (100 mg) and N-ethyl-N,N-diisopropylamine (1.5 eq) in dioxane (2 ml), was added 2-thiophenecarbonyl chloride (1.5 eq). The mixture was stirred overnight at room temperature. After addition of dichloromethane, the organic layer was washed with water and a sodium hydroxide solution and then dried over sodium sulfate. The solvent was removed to give an amorphous material, which was crystallized in diethylether to give the title compound (80 mg).

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.45 (d, 1H), 7.3-7.2 (m, 4H), 7.05 (t, 1H), 6.9 (m, 3H), 6.7 (m, 2H), 6.5 (d, 1H), 6.2 (t, 1H), 4.0 (m, 1H), 3.5 (m, 1H), 2.2 (m, 1H), 2.0 (m, 1H), 1.85 (s, 3H).

HPLC:  $r_T$ : 10.02, 97.8% purity

MS positive ESI: m/z (m+H)<sup>+</sup> = 377

### Example 100

#### Cis-N-(1-Benzoyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide

[0187] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: phenyl, R<sup>4</sup>: H, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol C, using appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 7.45 (d, 1H), 7.3-7.2 (m, 4H), 7.1 (t, 2H), 7.0 (t, 1H), 6.9 (m, 4H), 6.8 (t, 1H), 6.4 (m, 1H), 6.3 (t, 1H), 4.1 (m, 1H), 3.4 (m, 1H), 2.2 (m, 1H), 2.0 (m, 1H), 1.85 (s, 3H).

HPLC:  $r_T$ : 10.12, 98.6% purity

MS positive ESI: m/z (m+H)<sup>+</sup> = 371

### Example 101

#### Cis-N-[2-Ethyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

[0188] R<sup>1</sup>: phenyl, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: pyridin-3-yl, R<sup>4</sup>: ethyl, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol C, using appropriate reactants.

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 8.5 (m, 2H), 7.4-7.1 (m, 8H), 7.0 (m, 1H), 6.9 (t, 1H), 6.4 (d, 1H), 5.5 (m, 1H), 4.6 (m, 1H), 2.3 (m, 1H), 2.0 (s, 3H), 1.6 (m, 1H), 1.3 (m, 2H), 0.7 (t, 3H).

HPLC:  $r_T$ : 8.63, 96.7% purity

MS positive ESI: m/z (m+H)<sup>+</sup> = 400

**Example 102****Cis-N-Ethyl-N-[2-Methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide**

**[0189]** R<sup>1</sup>: CH<sub>2</sub>-CH<sub>3</sub>, R<sup>2</sup>: CH<sub>3</sub>, R<sup>3</sup>: pyridin-3-yl, R<sup>4</sup>: CH<sub>3</sub>, R<sup>5</sup>: H, R<sup>6</sup>: H, R<sup>7</sup>: H, R<sup>8</sup>: H.

The titled compound was prepared according to protocol C, using appropriate reactants. (0.046g, 60% yield)

<sup>1</sup>H NMR [CDCl<sub>3</sub>]: δ 8.5 (m, 2H), 7.3 (m, 1H), 7.2 (m, 1H), 7.1 (m, 1H), 7.0 (m, 1H), 6.9 (m, 1H), 6.5(m,1H), 5.9 (m, 0.5H), 4.9 (m, 1.5H), 4.0(m,0.2H), 3.6 (m, 0.4H), 3.2(m, 0.4H), 2.8 (m, 1H), 2.6(m, 0.5H), 2.3(m, 1.5H), 2.2 (m, 1.5H), 1.7(m, 0.5H), 1.4(m, 3H), 1.2(m, 3H).

HPLC: r<sub>T</sub>: 5.87, 100% purity

MS positive ESI: m/z (m+H)<sup>+</sup>= 338

**Biological results****Binding test**


**[0190]** The binding of CRTH2 antagonists was measured using membranes prepared from CRTH2-expressing L1.2 cells. To make the membrane preparation, 10 grams of cell pellet were resuspended in 20 ml of cold 50 mM Tris HCl (pH 7.7), and homogenized three times (10 seconds each) on ice with a Polytron tissue homogenizer. The suspension was then diluted to 80 ml with the same buffer, and centrifuged at 40,000 x g for 30 min. After discarding the supernatant, the pellet was washed once with cold 80 mM Tris HCl (pH 7.7) by repeating the resuspension, homogenization, and centrifugation described above. Finally, the pellet was resuspended in 20 ml of cold 50 mM Tris HCl (pH 7.7), aliquoted, and stored at - 80°C. On the day of the assay, the membranes were thawed on ice, and 25 µl was placed into each well of a 96-well microtiter plate. 5 µl of 0.5 M MgCl<sub>2</sub> and 10 µl of PVT-WGA scintillation proximity assay (SPA) beads (Amersham) were then added to each well and mixed thoroughly. 5 µl of the compounds to be tested were then added, followed by 5 µl of 30 nM <sup>3</sup>H-PGD<sub>2</sub> (200 Ci/mmol from NEN diluted 1:17) to initiate the binding reaction. The plate is then sealed and incubated at room temperature for at least 1 hour before being counted on a Wallac Microbeta Trilux scintillation counter. Non-specific binding is measured in wells containing 100 µM unlabeled PGD<sub>2</sub>. Active compounds are able to compete with <sup>3</sup>H-PGD<sub>2</sub> for binding to CRTH2 and are identified by a decrease in the number of cpm bound. The compounds of the invention were tested in the above assay and were found to be active CRTH2 antagonists. The IC<sub>50</sub> values of compounds of examples 1 to 102 are below 5µM.

**Actin polymerization test**

**[0191]** Actin polymerization was assayed using an actin-specific fluorescent label, phalloidin —fluorescein isocyanate conjugate, which binds polymerized actin fiber. This test can be performed using cells such as isolated human eosinophils or isolated human TH2-cells. The cells (preparations were resuspended at 1.1 x 10<sup>6</sup> cells/ml in RPMI 1640+ pen/strep +0.5 %FCS. The cell suspension was aliquoted (450µl /tube) into 5 ml tube. The cells were incubated 2 min at 37°C with the cells and then the appropriate stimulus (PGD<sub>2</sub>) was added followed exactly 25 seconds later by 500 µL of stopping solution, which contains lysophosphatidylcholine (0.5mg/ml), PBS, 6 to 8 % formaldehyde, and 0.2µM NBD-phalloidin in ME OH. The tube was allowed to sit at 4°C for an hour. The tube was then centrifuged at 1400 rpm for 5 minutes. The cell pellets were washed twice and then resuspended in 500 µl PBS. Each sample was then read on a FACS Caliber instrument. Cells were gated using the forward scatter/ side scatter data in the lymphocyte area. Responses were measured by the change in median FL-1 fluorescence between vehicle treated cells and stimulus treated cells. Test compounds were assayed in the presence and absence of PGD<sub>2</sub>, and compared to a sample containing PGD<sub>2</sub> alone. A compound that reduces PGD<sub>2</sub>-induced actin polymerization of CRTH2 cells is identified as a candidate CRTH2 antagonist.

**Claims**

1. A compound of general formula (I):



wherein,

- $R^1$  represents H,  $(C_1-C_4)$ alkyl,  $(C_2-C_4)$ alkenyl,  $(C_2-C_4)$ alkynyl or  $(CH_2)_m-R^1$ , in which

$R^1$  is selected from aromatic heterocycle, phenyl and  $(C_3-C_6)$ cycloalkyl

wherein the phenyl, the heterocycle and the cycloalkyl groups are unsubstituted or substituted by one or several groups selected from

- $Q^1$ , and,
- $(C_1-C_4)$ alkyl optionally substituted with one or several groups which are the same or different and which are selected from  $Q^1$ ,

wherein  $Q^1$  is selected from halogen,  $NO_2$ , CN,  $SO_2CH_3$ ,  $SO_2NR^9R^{10}$ ,  $OR^9$ ,  $COOR^9$ ,  $C(=O)NR^9R^{10}$ ,  $NR^9R^{10}$ ,  $NR^9SO_2R^{10}$ ,  $NR^9C(=O)R^{10}$  and  $C(=O)R^9$  wherein  $R^9$  and  $R^{10}$  are the same or different and are selected from H and  $(C_1-C_4)$ alkyl;

$m$  is an integer selected from 0, 1 and 2;

- $R^2$  represents  $(C_1-C_4)$ alkyl, wherein the alkyl group may be substituted with one or several, substituents selected from halogen,  $OR^9$ ,  $NR^9R^{10}$ ,  $COOR^9$ ,  $C(=O)NR^9R^{10}$ ,  $NHSO_2R^9$  and  $C(=O)(C_1-C_4)$ alkyl wherein  $R^9$  and  $R^{10}$  are the same or different and are selected from H and  $(C_1-C_4)$ alkyl;
- $R^3$  represents  $(C_3-C_6)$ cycloalkyl or  $-A-R^3$ , wherein
- $A$  represents a bond, straight or branched  $(C_1-C_3)$ alkylene, or  $(C_2-C_3)$ alkenylene;
- $R^3$  represents  $(C_6-C_{12})$ aryl or an heterocycle, optionally aromatic, having from 5 to 10 atoms in the cycle, wherein the aryl and the heterocycle groups are unsubstituted or substituted by one or several substituents selected from,
- an aryle, an aromatic heterocycle,
- $Q^2$ , and,
- $(C_1-C_4)$ alkyl optionally substituted with one or several groups which are the same or different and which are selected from  $Q^2$ ,

wherein  $Q^2$  is selected from halogen,  $NO_2$ , CN,  $SO_2CH_3$ ,  $SO_2NR^9R^{10}$ ,  $OR^9$ ,  $SR^9$ ,  $COOR^9$ ,  $C(=O)NR^9R^{10}$ ,  $NR^9R^{10}$ ,  $NR^9SO_2R^{10}$ ,  $NR^9C(=O)R^{10}$  and  $C(=O)R^9$  wherein  $R^9$  and  $R^{10}$  are the same or different and are selected from H and  $(C_1-C_4)$ alkyl;

- $R^4$  represents H or  $(C_1-C_4)$ alkyl;


- $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are the same or different and are selected from

- H,  $Q^3$ , and,
- $(C_1-C_4)$ alkyl optionally substituted with one or several groups which are the same or different and which

are selected from Q<sup>3</sup>,

wherein Q<sup>3</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, SR<sup>9</sup> COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl; their optical isomers, as well as their N-oxides and pharmaceutically acceptable salts, as a medicament.

## 2. A compound of formula (I)





wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in claim 1, with the exclusion of the following compounds;

N-(1-benzoyl-6-bromo-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-acetanilide,  
 N-(1-benzoyl-6-chloro-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-acetanilide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-(4-methoxyphenyl)-2-methyl-propanamide,  
 N-[1-(4-fluorobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-butanamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-pentanamide,  
 N-[1-[(4-fluorophenyl)acetyl]-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-propanamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-2,2-dimethyl-N-phenyl-propanamide,  
 N-(1-benzoyl-6-bromo-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-phenyl-pentanamide,  
 N-[1-(2-furanylcabonyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 2-methyl-N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-propanamide,  
 2,2,2-trifluoro-N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-acetamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-(3-methoxyphenyl)-acetamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-(4-methylphenyl)-acetamide,  
 N-[1-(4-chloro-3-nitrobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(3-nitrobenzoyl)-4-quinolyl]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-[3-(4-methoxyphenyl)-1-oxo-2-propenyl]-2-methyl-4-quinolyl]-acetamide,  
 N-[1-(3-chlorobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 N-[1-(3-fluorobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 N-[1-[4-(1,1-dimethylethyl)benzoyl]-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenyl-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(1-oxo-3-phenyl-2-propenyl)-4-quinolyl]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(2-thienylcarbonyl)-4-quinolyl]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-(4-methoxybenzoyl)-2-methyl-4-quinolyl]-acetamide,  
 N-[1-(3,5-dinitrobenzoyl)-1,2,3,4-tetrahydro-2-methyl-4-quinolyl]-N-phenylacetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-2-methyl-1-(4-nitrobenzoyl)-4-quinolyl]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-(2-iodobenzoyl)-2-methyl-4-quinolyl]-acetamide,  
 N-phenyl-N-[1,2,3,4-tetrahydro-1-(3-methoxybenzoyl)-2-methyl-4-quinolyl]-acetamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-phenyl-pentanamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-phenyl-butanamide,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinolyl)-N-phenyl-propanamide, 1-benzoyl-1,2,3,4-tetrahydro-4-(N-phenylacetamido)-quinaldine,

N-[(1-benzoyl-1, 2, 3, 4-tetrahydro-2-methyl-4-quinoliny)]-2-methyl-N-phenyl propanamide;  
 N-[1-(4-bromobenzoyl)-1,2,3,4-tetrahydro-2,6-dimethyl-4-quinoliny]]-acetamide;  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2,6-dimethyl-4-quinoliny)]-acetamide; and,  
 N-(1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinoliny)]-acetamide,

its optical isomers, as well as its N-oxides and pharmaceutically acceptable salts.

3. A compound of formula (Ia) or (Ib), or a mixture thereof,





wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in claim 1 and R<sup>4</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl, with the exclusion of the following compounds,

N-[(2R, 4S)-1-benzoyl-1, 2, 3, 4-tetrahydro-2-methyl-4-quinoliny]]-2-methyl-N-phenyl propanamide;  
 N-[(2R,4S)-1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinoliny]]-2,2-dimethyl-N-phenyl-propanamide;  
 N-[(2R, 4S)-1-benzoyl-1, 2, 3, 4-tetrahydro-2-methyl-4-quinoliny]]-N-phenyl butanamide;  
 N-[(2R, 4S)-1-benzoyl-1, 2, 3, 4-tetrahydro-2-methyl-4-quinoliny]]-N-phenyl acetamide;  
 N-[(2R,4S)-1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinoliny]]-N-phenyl-pentanamide,  
 N-[(2R,4S)-1-benzoyl-1,2,3,4-tetrahydro-2-methyl-4-quinoliny]]-acetamide,

its optical isomers, as well as its N-oxides and pharmaceutically acceptable salts.

4. A compound of formula (Ic) or (Id), or a mixture thereof



wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in claim 1 and R<sup>4</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl.

5. A compound according to any one of claims 2, 3 and 4 wherein R<sup>1</sup> represents H, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>2</sub>-C<sub>4</sub>)alkenyl, (C<sub>2</sub>-C<sub>4</sub>)alkynyl or (CH<sub>2</sub>)<sub>m</sub>-R<sup>1</sup>, wherein

R<sup>1</sup> is selected from phenyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl wherein the phenyl and the cycloalkyl groups are unsubstituted or substituted by one or several groups selected from

- Q<sup>1</sup>, and,
- (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several groups which are the same or different and which are selected from Q<sup>1</sup>,

wherein Q<sup>1</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;

m is an integer selected from 0, 1 and 2.

6. A compound according to claim 5 wherein R<sup>1</sup> represents (CH<sub>2</sub>)<sub>m</sub>-R<sup>1</sup>, wherein

R<sup>1</sup> is selected from phenyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl wherein the phenyl and the cycloalkyl groups are unsubstituted or substituted by 1 to 3 groups selected from OR<sup>9</sup>, COOR<sup>9</sup> and (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with a group COOR<sup>9</sup>, and

m is an integer selected from 0 and 1.

7. A compound according to any one of claims 2, 3 and 4 wherein R<sup>1</sup> represents (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>2</sub>-C<sub>4</sub>)alkenyl, (C<sub>2</sub>-C<sub>4</sub>)alkynyl or (CH<sub>2</sub>)<sub>m</sub>-R<sup>1</sup>, in which

R<sup>1</sup> is selected from aromatic heterocycle and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl wherein the heterocycle and the cycloalkyl groups are unsubstituted or substituted by one or several groups selected from

- Q<sup>1</sup>, and,
- (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several groups which are the same or different and which are selected from Q<sup>1</sup>,

wherein Q<sup>1</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl;

m is an integer selected from 0, 1 and 2.

8. A compound according to claim 7 wherein R<sup>1</sup> represents a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl wherein the cycloalkyl group is unsubstituted or substituted by one or several groups selected from

- Q<sup>1</sup>, and,
- (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several groups which are the same or different and which are selected from Q<sup>1</sup>,

wherein Q<sup>1</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl.

9. A compound according to claim 8 wherein R<sup>1</sup> represents a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl.

10. A compound according to claim 5 wherein R<sup>1</sup> represents phenyl unsubstituted or substituted in the para position by a substituent selected from halogen, OR<sup>9</sup>, CH<sub>2</sub>COOR<sup>9</sup> and CH<sub>2</sub>COOR<sup>9</sup> wherein R<sup>9</sup> is selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl.

11. A compound according to any one of claim 2 to 10 wherein R<sup>2</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl.

12. A compound according to any one of claim 2 to 10 wherein R<sup>4</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl.

13. A compound according to any one of claims 2 to 12 wherein R<sup>3</sup> is selected from (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl and -A-R<sup>3</sup>, wherein

- A represents a bond, (C<sub>1</sub>-C<sub>3</sub>)alkylene, straight or branched, or (C<sub>2</sub>-C<sub>3</sub>)alkenylene;
- R<sup>3</sup> represents a heterocycle, optionally aromatic, having from 6 to 10 atoms in the cycle, unsubstituted or substituted by one or several, preferably 1 to 3, substituents selected from,
- an aryle, an heterocycle,
- Q<sup>2</sup>, and,
- (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several, preferably 1 to 3, groups which are the same or different and which are selected from Q<sup>2</sup>,

wherein Q<sup>2</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, SR<sup>9</sup>COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl, with the proviso that R<sup>3</sup> is not selected from unsubstituted thienyl or unsubstituted furanyl.

14. A compound according to any one of claims 2 to 12 wherein R<sup>3</sup> is selected from -A-R<sup>3</sup>, wherein

- A represents a bond, straight or branched (C<sub>1</sub>-C<sub>3</sub>)alkylene, or (C<sub>2</sub>-C<sub>3</sub>)alkenylene;
- R<sup>3</sup> represents a phenyl, unsubstituted or substituted by one or several substituents selected from,
- an aryle, an heterocycle,
- Q<sup>2</sup>, and,
- (C<sub>1</sub>-C<sub>4</sub>)alkyl optionally substituted with one or several groups which are the same or different and which are selected from Q<sup>2</sup>,

wherein Q<sup>2</sup> is selected from halogen, NO<sub>2</sub>, CN, SO<sub>2</sub>CH<sub>3</sub>, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, OR<sup>9</sup>, SR<sup>9</sup>COOR<sup>9</sup>, C(=O)NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>R<sup>10</sup>, NR<sup>9</sup>SO<sub>2</sub>R<sup>10</sup>, NR<sup>9</sup>C(=O)R<sup>10</sup> and C(=O)R<sup>9</sup> wherein R<sup>9</sup> and R<sup>10</sup> are the same or different and are selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl.

15. A compound according to any one of claims 2 to 14 wherein R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are the same or different and are selected from H, halogen and OR<sup>9</sup> wherein R<sup>9</sup> is selected from H and (C<sub>1</sub>-C<sub>4</sub>)alkyl.

16. A compound selected from the group consisting of

Cis-N-[2-Methyl-1-(pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(1-oxy-pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(4-Hydroxy-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(4-trifluoromethyl-benzoyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(4-Cyano-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(4-Chloro-benzoyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 4-[Cis-4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-benzoic acid methyl ester;  
 4-[Cis-4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-benzoic acid;  
 Cis-N-[2-Methyl-1-(3-phenyl-propionyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(5-methyl-thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(Benzofurazan-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-(2-Methyl-1-phenylacetyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(pyrazine-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(6-Chloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(6-trifluoromethyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2,6-Dimethoxy-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2-Methoxy-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(2-methylsulfanyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[1-(2-Chloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-[2-Methyl-1-(5-methyl-pyrazine-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;



Cis-N-[1-(2-Chloro-6-methyl-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[1-(4-Chloro-1,3-dimethyl-1H-pyrazolo[3,4-b]pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[2-Methyl-1-(6-(2,2,2-trifluoro-ethoxy)-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[2-Methyl-1-(2-propylsulfanyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[1-(5,6-Dichloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[1-(2,6-Dichloro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[2-Methyl-1-(4-methyl-[1,2,3]thiadiazole-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[2-Methyl-1-(5-methyl-isoxazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[1-(2,5-Dimethyl-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[2-Methyl-1-(1-methyl-1H-pyrrole-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[1-(Isoxazole-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[2-Methyl-1-(5-methyl-isoxazole-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[1-(2,4-Dimethyl-thiazole-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[1-(5-Chloro-thiophene-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[1-(1,5-Dimethyl-1H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;

Cis-N-[2-Methyl-1-(4-methyl-isothiazole-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-5-[4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-thiophene-2-carboxylic acid dimethylamide

Cis-N-[1-(4-Hydroxy-quinoline-6-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(4-Hydroxy-quinoline-6-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(4-tert-Butyl-thiazole-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(2-Ethyl-pyridine-4-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(3,6-Dichloro-pyridine-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(4-Chloro-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-2-[4-(Acetyl-phenyl-amino)-2-methyl-3,4-dihydro-2H-quinoline-1-carbonyl]-isonicotinic acid methyl ester

Cis-N-[2-Methyl-1-(4-[1,2,4]triazol-4-yl-benzoyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(2,6-Dimethoxy-pyridine-4-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(5-Ethyl-isoxazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[2-Methyl-1-(2-tetrazol-1-yl-pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[2-Methyl-1-(5-propyl-isoxazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(5-Isobutyl-2-methyl-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(5-Bromo-furan-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[2-Methyl-1-(6-phenyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[2-Methyl-1-(2-phenyl-pyridine-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[2-Methyl-1-(quinoline-6-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(3,4-Dimethoxy-furan-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[2-Methyl-1-(3-methyl-furan-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(2,5-Dimethyl-furan-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(2,4-Dimethyl-oxazole-5-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(5-Methoxymethyl-furan-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(5-Fluoro-pyridine-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[2-Methyl-1-(quinoline-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[2-Methyl-1-(6-methyl-pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[2-Methyl-1-(quinoline-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[2-Methyl-1-(1H-pyrazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[2-Methyl-1-(2H-pyrazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[1-(5-Isobutyl-isoxazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[2-Methyl-1-(quinoline-4-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[2-Methyl-1-(6-methyl-pyridine-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide

Cis-N-[2-Methyl-1-(quinoxaline-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(3-Methoxy-thiophene-2-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[1-(5-tert-Butyl-2-methyl-furan-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 5 Cis-N-[1-(5-Ethyl-2-methyl-2H-pyrazole-3-carbonyl)-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-([1,2,5]thiadiazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 Cis-N-[2-Methyl-1-(2-methyl-5-propyl-2H-pyrazole-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide  
 10 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-Benzylacetamide;  
 Cis-N-Benzyl-N-[2-methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Trans-N-Benzyl-N-[2-methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-Cyclohexyl-N-[2-methyl-1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-(1-Benzoyl-6-methoxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 15 Cis-N-(1-Benzoyl-6-hydroxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 Cis-N-(1-Benzoyl-6-chloro-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide;  
 N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-prop-2-ynyl-acetamide;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-(4-methoxy-phenyl)-acetamide;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-(4-hydroxyphenyl)-acetamide;  
 20 Cis-[4-[Acetyl-(1-benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-amino]-phenyl]-acetic acid ethyl ester;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-malonamic acid methyl ester;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-malonamic acid;  
 Cis-N-[2-Methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide;  
 Cis-N-(1-Benzoyl-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-Cyclopropyl-acetamide;  
 25 Cis-N-Cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide  
 (+)-Cis-N-cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide  
 (-)-Cis-N-cyclopropyl-N-[2-methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-Cyclopropyl-N-[2-methyl-1-(3-methyl-isoxazole-5-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 30 Cis-N-Phenyl-N-[1-(thiophene-2-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide;  
 Cis-N-(1-Benzoyl-1,2,3,4-tetrahydro-quinolin-4-yl)-N-phenyl-acetamide  
 Cis-N-[2-Ethyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-N-phenyl-acetamide and  
 Cis-N-Ethyl-N-[2-Methyl-1-(pyridine-3-carbonyl)-1,2,3,4-tetrahydro-quinolin-4-yl]-acetamide.

35 17. A compound according to any one of claims 2 to 16 as a medicament.

18. A pharmaceutical composition comprising a compound of formula (I) according to any one of claims 1 to 16 together with a pharmaceutically acceptable carrier, excipient, diluent or delivery system.

40 19. Use of a compound of formula (I) according to any one of claims 1 to 16 for the preparation of a medicament for the prevention or the treatment of disorders for which therapy by a CRTH2 antagonist is relevant.

20. Use according to claim 19 in which said disorder is selected from inflammatory disorders such as rheumatoid arthritis, osteoarthritis, atherosclerosis, Crohn's disease, colitis ulcerosa, inflammatory bowel disease; disorders of the skin including psoriasis, eczema, erythema, pruritis, and acne; systemic lupus erythematosus, chronic obstructive pulmonary disease, angioedema, stroke, and any disease marked by reperfusion injury, graft rejection, and autoimmune diseases, allergic diseases, such as allergic asthma, atopic dermatitis, and allergic rhinitis.



21. Use according to claim 20 in which said disorder is selected from asthma and allergic rhinitis.

22. Method for the treatment of a disorder for which therapy by a CRTH2 antagonist is relevant, comprising administering to a mammal in need thereof an effective amount of compound of formula (I) according to any one of claims 1 to 16.

23. Method according to claim 22 in which said disorder is selected from inflammatory disorders such as rheumatoid arthritis, osteoarthritis, atherosclerosis, Crohn's disease, colitis ulcerosa, inflammatory bowel disease; disorders of the skin including psoriasis, eczema, erythema, pruritis, and acne; systemic lupus erythematosus, chronic obstructive pulmonary disease, angioedema, stroke, and any disease marked by reperfusion injury, graft rejection,


and autoimmune diseases, allergic diseases, such as allergic asthma, atopic dermatitis, and allergic rhinitis.

24. A method for the preparation of a compound of formula (Ia) or (Ib) or mixture thereof,



said method comprising the following steps;


(1) reacting a compound of formula (3a)



wherein R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in claim 1, with


- a compound of formula R<sup>4</sup>-MgCl, wherein R<sup>4</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl in an anhydrous ether such as tetrahydrofuran, and then with,
- a compound of formula R<sup>3</sup>-COCl,

to give a compound of formula (3b)




wherein R<sup>3</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in claim 1 and R<sup>4</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl;

(2) reacting said compound of formula (3b) with a compound of formula  $R^1-NH_2$ , wherein  $R^1$  is as defined in claim 1, in the presence of a base, to give a compound of formula (3c),



wherein  $R^1$ ,  $R^3$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined in claim 1 and  $R^4$  is  $(C_1-C_4)$ alkyl;

(3) reducing said compound of formula (3c) with a mineral borohydride such as sodium triacetoxo borohydride in acetic acid to give a compound of formula (3d),





wherein  $R^1$ ,  $R^3$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined in claim 1 and  $R^4$  is  $(C_1-C_4)$ alkyl;

(4) reacting said compound of formula (3d) with a compound of formula  $R^2COCl$ , wherein  $R^2$  is as defined in claim 1, in the presence of a base, to give a mixture of compounds of formula (Ia) and (Ib) wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined in claim 1 and  $R^4$  is  $(C_1-C_4)$ alkyl;

(5) isolating said mixture, or


(6) separating and isolating said compounds of formula (Ia) and (Ib).

25. A method for the preparation of a compound of formula (Ia) or (Ib) or a mixture thereof,




said method comprising the following steps,

(1) reacting a compound of formula (4a)



(4a)


with a compound of formula  $R^1-NH_2$ , wherein  $R^1$  is as defined in claim 1, to give the compound of formula (4b),



(4b)

in which  $R^1$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined in claim 1;


(2) reacting said compound of formula (4b), with a compound of formula  $R^2COCl$ , wherein  $R^2$  is as defined in claim 1, in the presence of a base, to give the compound of formula (4c),



(4c)


in which  $R^1$ ,  $R^2$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined in claim 1;

(3) reacting said compound of formula (4c) with benzyl-halogen in a solvent such as acetone to give the compound of formula (4d),




in which  $R^1$ ,  $R^2$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined in claim 1;

(4) optionally, reacting said compound of formula (4d) with a compound of formula  $R^4\text{-MgCl}$ , wherein  $R^4$  is  $(C_1\text{-}C_4)$ alkyl in an anhydrous ether such as tetrahydrofuran to give the compound of formula (4e),




in which  $R^1$ ,  $R^2$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are as defined in claim 1 and  $R^4$  is  $(C_1\text{-}C_4)$  alkyl;

(5) reducing said compound of formula (4d) or (4e), for example by hydrogenation in MeOH in presence (R, R)-(-)-1,2-bis[(o-methoxyphenyl)(phenyl)phosphino]ethane (1,5-cyclooctadiene)rhodium (I) tetrafluoroborate or  $\text{NiCl}_2/\text{NaBH}_4$ , to give a compound of formula (4f)



20 in which R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in claim 1 and R<sup>4</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>) alkyl;

(6) deprotecting said compound of formula (4f) by hydrogenolysis using for example ammonium formate catalysed by palladium on charcoal, to give a compound of formula (4g)



40 in which R<sup>1</sup>, R<sup>2</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined in claim 1;

(7) reacting said compound of formula (4g) with R<sup>3</sup>COCl, wherein R<sup>3</sup> is as defined in claim 1, in the presence of a base, to give a mixture compounds of formula (Ia) and (Ib) wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined above;

45 (8) isolating said compounds of formula (Ia) and (Ib), or,

(9) separating compounds of formula (Ia) and (Ib) and isolating them.

50

55



European Patent  
Office

# PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 03 29 0025  
shall be considered, for the purposes of subsequent  
proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	WO 00 17165 A (WESTER RONALD THURE ; PFIZER PROD INC (US); DENINNO MICHAEL PAUL (U) 30 March 2000 (2000-03-30) * claims 1,33 *	1,18,20, 24,25	C07D409/06 C07D215/44 C07D401/06 C07D413/06 C07D471/04
D,X	WO 02 079165 A (NIKITIDIS GRIGORIOS ; BLADH HAKAN (SE); NORDEN CARITA (SE); ASTRAZE) 10 October 2002 (2002-10-10) * examples 8,14; table I * * claims 2-14,17,20; table II * * page 22 * --- -/--	1,18,20, 24,25	C07D417/06 C07D401/10 C07D405/06 C07D401/14 A61K31/4704 A61K31/4709 A61P9/10 A61P17/06 A61P29/00 A61P37/00
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			C07D A61K A61P
<b>INCOMPLETE SEARCH</b>			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search:</p> <p>Although claims 22 and 23 are directed to a method of treatment of the human/animal body (Article 52(4) EPC), the search has been carried out and based on the alleged effects of the compound/composition.</p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		18 June 2003	Seitner, I
CATEGORY OF CITED DOCUMENTS		<p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>	
<p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p>			

EPO FORM 1503 03/02 (P04C07)





European Patent  
Office

## PARTIAL EUROPEAN SEARCH REPORT

Application Number  
EP 03 29 0025

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	<p>DATABASE CAPLUS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; ZALUKAEV, L. P. ET AL: "Intramolecular donor-acceptor interaction in 2-7ethyl-4-anilino-1,2,3,4-tetrahydroquinoline and its derivatives" retrieved from STN Database accession no. 69:27206 XP002244839 * CAS RN: 13125-49-0; 17117-38-3 * &amp; TRUDY PROBLEMNOI LABORATORII KHIMII VYSOKOMOLEKULYARNYKH SOEDINENII, VORONEZHSKII GOSUDARSTVENNYI UNIVERSITET (1966), NO. 4, 5-16 ,</p>	1	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
X	<p>--- DATABASE CAPLUS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; ZALUKAJEVS, L. ET AL: "Bimolecular alkylidenearylamines. VIII. Synthesis and bromination of 2-methyl-4-N-acetylanilino-1,2,3,4-tetrahydroquinoline" retrieved from STN Database accession no. 59:54789 XP002244840 * CAS RN: 95868-01-2 * &amp; ZHURNAL OBSHCHEI KHIMII (1963), 33(6), 1956-8 ,</p> <p>--- -/--</p>	1	

EPO FORM 1503 03.82 (P04C10)



European Patent  
Office

## PARTIAL EUROPEAN SEARCH REPORT

Application Number  
EP 03 29 0025

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	<p>DATABASE CHEMCATS [Online] CHEMICAL ABSTRACT SERVICE, COLUMBUS, OHIO, US; retrieved from STN XP002244841 * Order Number: MCL016056 * &amp; "MCL Screening Compounds" 24 May 2001 (2001-05-24) , MCL MOSCOW LTD. , MOSCOW, 121170, RUSSIA ---</p>	1	
X	<p>DATABASE CHEMCATS [Online] CHEMICAL ABSTRACT SERVICE, COLUMBUS, OHIO, US; retrieved from STN XP002244842 * Order Number: C-013353; K-021540; X-077207; X-015189; POD_54/0172; POD_35/0696 * &amp; "Scientific Exchange Product List" 1 January 2003 (2003-01-01) , SCIENTIFIC EXCHANGE, INC. , CENTER OSSIPPEE, NH, 03814, USA ---</p>	1	<p>TECHNICAL FIELDS SEARCHED (Int.Cl.7)</p>
X	<p>DATABASE CHEMCATS [Online] CHEMICAL ABSTRACT SERVICE, COLUMBUS, OHIO, US; retrieved from STN XP002244843 * Order Number: T0501-8575; T0400-0627; T0202-0105; T0200-3749 * &amp; "Enamine Product Listing" 15 November 2001 (2001-11-15) , ENAMINE , KIEV 042, 01042, UKRAINE ---</p> <p style="text-align: center;">-/--</p>	1	



European Patent  
Office

## PARTIAL EUROPEAN SEARCH REPORT

Application Number  
EP 03 29 0025

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	DATABASE CHEMCATS [Online] CHEMICAL ABSTRACT SERVICE, COLUMBUS, OHIO, US; retrieved from STN XP002244844 * Order Number: 17512; 16101; 15586; 15485; 15102; 15101 * & "MicroChemistry Screening Collection" 3 October 2002 (2002-10-03) , MICROCHEMISTRY-RADAPHARMA , MOSCOW, 111123, RUSSIA ---	1	
X	DATABASE CHEMCATS [Online] CHEMICAL ABSTRACT SERVICE, COLUMBUS, OHIO, US; retrieved from STN XP002244845 * Order Number: VIT015003; VIT009313; VIT007916; VIT004828; VIT016130; VIT017578 * & "Vitas-M Screening Collection" 22 March 2001 (2001-03-22) , VITAS-M , MOSCOW, 1192829, RUSSIA ---	1	TECHNICAL FIELDS SEARCHED (Int.Cl.7)
X	DATABASE CHEMCATS [Online] CHEMICAL ABSTRACT SERVICE, COLUMBUS, OHIO, US; retrieved from STN XP002244846 * Order Number: 6101326; 5805939; 5481700; 5304469; 5228582; 5228571 * & "ChemBridge Product List" 17 January 2002 (2002-01-17) , CHEMBRIDGE CORPORATION , SAN DIEGO, CA, 92127, USA --- -/--	1	

EPO FORM 1503 03.02 (P04C10)



European Patent  
Office

## PARTIAL EUROPEAN SEARCH REPORT

Application Number  
EP 03 29 0025

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	<p>DATABASE CHEMCATS [Online]            CHEMICAL ABSTRACT SERVICE, COLUMBUS, OHIO, US;            retrieved from STN            XP002244847            * Order Number: AK-968/37032032;            STOCK1S-91726; STOCK1S-91121;            STOCK1S-63562; STOCK1S-59611;            STOCK1S-57645; STOCK1S-54769;            STOCK1S-53637; STOCK1S-04860; 3535-0169;            3390-0230; BAS 2927251            BAS 1921878; BAS 0916297; BAS 0757452; BAS            0757450; BAS 0458946; BAS 0315505; BAS            0225055; BAS 0126897; BAS 0126850; BAS            0107978 *            &amp; "Interchim Intermediates"            9 July 2002 (2002-07-09) , INTERCHIM ,            MONTLUCON, CEDEX, 03103, FRANCE            ---</p>	1	
D,A	<p>EP 0 987 251 A (PFIZER PROD INC)            22 March 2000 (2000-03-22)            * claims 1,30,40 *            -----</p>	1,18,20	<p>TECHNICAL FIELDS            SEARCHED (Int.Cl.7)</p>

EPO FORM 1503 03.02 (P04C10)

ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.

EP 03 29 0025

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

18-06-2003

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0017165 A	30-03-2000	AU 747715 B2	23-05-2002
		AU 5440399 A	10-04-2000
		BG 105429 A	31-12-2001
		BR 9913855 A	24-07-2001
		CA 2344248 A1	30-03-2000
		CN 1318057 T	17-10-2001
		CZ 20010954 A3	12-12-2001
		EE 200100167 A	17-06-2002
		EP 1114032 A1	11-07-2001
		HR 20010200 A1	30-04-2002
		HU 0103512 A2	28-02-2002
		WO 0017165 A1	30-03-2000
		JP 2002526476 T	20-08-2002
		NO 20011349 A	14-05-2001
		NZ 509837 A	26-11-2002
		PL 346830 A1	25-02-2002
		SK 3532001 A3	03-12-2001
		TR 200100780 T2	23-07-2001
		US 6140343 A	31-10-2000
		US 6489478 B1	03-12-2002
		ZA 200101745 A	02-05-2002
WO 02079165 A	10-10-2002	WO 02079165 A1	10-10-2002
EP 0987251 A	22-03-2000	BR 9904141 A	12-09-2000
		EP 0987251 A1	22-03-2000
		JP 2000191645 A	11-07-2000
		US 6140342 A	31-10-2000
		US 6362198 B1	26-03-2002